

Recombinant full-length mouse CAMK1b kinase containing N-terminal GST tag

Catalog No.	CRC105A CRC105B	Quantity:	5 µg 20 µg
Description:	<p>Recombinant full-length mouse CAMK1b containing N-terminal GST tag was expressed by baculovirus in Sf 9 insect cells.</p> <p>Many of the effects of calcium are mediated via its interaction with calmodulin and the subsequent activation of Ca(2+)/calmodulin-dependent (CaM) kinases. CaM kinases are involved in a wide variety of cellular processes including muscle contraction, neurotransmitter release, cell cycle control, and transcriptional regulation. While CaMKII has been implicated in learning and memory, the biological role of the other multifunctional CaM kinases, CaMKI and CaMKIV, is largely unknown. CaMKIbeta, or pregnancy upregulated non-ubiquitously expressed CaM kinase (PNCK), is a 38-kDa serine/threonine kinase whose catalytic domain shares 45-70% identity with members of the CaM kinase family. The gene for CaMKIbeta localizes to mouse chromosome X. CaMKIbeta is upregulated during intermediate and late stages of murine fetal development with highest levels of expression in developing brain, bone, and gut. CaMKIbeta is also expressed in a tissue-specific manner in adult mice with highest levels of expression detected in brain, uterus, ovary, and testis. Interestingly, CaMKIbeta expression in these tissues is restricted to particular compartments and appears to be further restricted to subsets of cells within those compartments. The chromosomal localization of CaMKIbeta, along with its tissue-specific and restricted pattern of spatial expression during development, suggests that CaMKIbeta may be involved in a variety of developmental processes including development of the central nervous system. Also CaMKIbeta2, an isoform of mCaMKIbeta, was mainly identified in the nervous system, including brain, spinal cord, trigeminal ganglion, and retina. Within the CNS, the expression of CaMKIbeta2 is detected in the mantle zone, but not in the ventricular zone, suggesting its possible involvement in the differentiation of neurons.</p>		
Concentration:	0.1 mg/ml		
Gene Accession No.:	NM_012040		
Source:	Sf9 insect cells		
Formulation:	Recombinant protein in storage buffer (50 mM Tris-HCl + 150 0.1 mM EGTA + 0.1 mM EDTA + 0.1 mM PMSF + 25% glycerol; pH 7.5).		
Purity:	2 µg of CAMK1b protein was subjected to SDS-PAGE and Coomassie blue staining. The scan of the gel showed >90% purity of the CAMK1b band product, and the band was at ~64 kDa (Fig. 2).		
Specific Activity:	223 nmol/min/mg: 223 nmol phosphate incorporated into Autocamtide 2 per minute per mg protein at 30°C for 15 minutes using a final concentration of 50 µM ATP (0.83 µCi/assay). See QA/QC section for details.		
Storage & Stability:	Store product frozen at or below -70°C. Stable for 1 year at -70°C as undiluted stock. Aliquot to avoid repeated thawing and freezing.		



QA/QC:

CAMK1b protein (100 ng/μl concentration) was diluted to 25 ng/μl in assay dilution buffer (4 mM MOPS + 2.5 mM β-glycerophosphate + 1 mM EGTA + 0.4 mM EDTA + 4 mM MgCl₂ + 0.05 mM DTT + 40 ng/μl BSA + 40 ng/μl BSA; pH 7.2), followed by 2-fold serial dilutions, and then the 10μl diluted proteins were used to phosphorylate the Autocamide 2 (KKALRRQETVDAL-amide) in the following assay condition:

- 10 μl diluted CAMK1b protein
- 7.5 μl Autocamide 2 (1 mg/ml stock)
- 2.5 μl Calmodulin (0.3 mg/ml in 5 mM CaCl₂)
- 5 μl [³²P] ATP mixture (250 μM ATP stock, 0.16 μCi/μl in 4x assay dilution buffer)

The various reaction components, except [³²P] ATP, were incubated at 30°C and the reaction started by the addition of [³²P] ATP. After 15 minutes, the reaction was terminated by spotting 20 μl of the reaction mixture onto a phosphocellulose P81 paper. The P81 paper was dried and washed several times in 1% phosphoric acid prior to counting in the presence of scintillation fluid in a scintillation counter. The actual counts, using various dilutions of the enzyme in the assay, are shown in Fig. 1.

Fig. 1 CAMK1b activity assay





Fig. 2 CAMK1b protein gel



Please note: always centrifuge vials before opening.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

