

NPB353Ge01 100µg
Advanced Glycation End Product (AGE)
Organism Species: General
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose.

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[RELEVANCE]

An advanced glycation end-product (AGE) is the result of a chain of chemical reactions after an initial glycation reaction. Glycation is accomplished by the Maillard reaction, which is a multistep process that begins with Schiff base formation between the amine and the carbonyl group on the sugar followed by rearrangement to form Amadori intermediates. AGEs affect nearly every type of cell and molecule in the body, and are thought to be one factor in aging and some age-related chronic diseases. BSA is glycosylated by the reaction of the glucose and BSA in vitro, and then being multistep purification, ending in high purified AGEs.

[**USAGE**]

Reconstitute in ddH₂O.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.