

RPA525Mu01 10 μ g

Recombinant Plasminogen Activator, Tissue (tPA)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[**PROPERTIES**]

Source: Prokaryotic expression

Host: *E.coli*

Residuess: Glu359~Leu532

Tags: N-terminal His Tag

Tissue Specificity: Secreted, Extracellular matrix

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.5

Predicted Molecular Mass: 20.7kDa

Accurate Molecular Mass: 23kDa as determined by SDS-PAGE reducing conditions.

[**USAGE**]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

ER FPPNHLKVVL GRTYRVVPG E EQTFEIEKY IVHEEFDDT
YDNDIALLQL RSQSKQCAQE SSSVGTACL DPNLQLPDWT ECELSGYGKH
EASSPFFSDR LKEAHVRLYP SSRCTSQH LF NKTVTNNMLC AGDTRSGGNQ
DLHDACQGDS GGPLVCMINK QMTLTGIISW GL

[**IDENTIFICATION**]




Figure. SDS-PAGE