

RPD471Hu01 100µg
Recombinant Pappalysin 2 (PAPPA2)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[**PROPERTIES**]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Gly888~Ala1093

Tags: N-terminal His-Tag

Tissue Specificity: Breast, Kidney, Brain, Pancreas.

Subcellular Location: Secreted.

Purity: >95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.5

Predicted Molecular Mass: 24.4kDa

Accurate Molecular Mass: 27kDa as determined by SDS-PAGE reducing conditions.

[**USAGE**]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

				GTF RQYVHTASSR
RVCDSSGYWT	PEEAVGPPDV	DQPCEPSLQA	WSPEVHLYHM	NMTVPCPTeg
CSLELLFQHP	VQADTLTLWV	TSFFMESSQV	LFDTEILLEN	KESVHLGPLD
TFCDIPLTIK	LHVDGKVSGV	KVYTFDERIE	IDAALLTSQP	HSPLCSGCRP
VRYQVLRDPP	FASGLPVVVT	HSHRKFTDVE	VTPGQMYQYQ	VLA

[IDENTIFICATION]




Figure 1. SDS-PAGE