

RPE275Hu01 50µg

Recombinant Spleen Tyrosine Kinase (SYK)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Asp374~Asn635

Tags: N-terminal His Tag

Subcellular Location: Membrane, Cytoplasm

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.4

Predicted Molecular Mass: 34.0kDa

Accurate Molecular Mass: 34kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

			DKELGSG	NFGTVKKGY	QMKKVVKTVA
VKILKNEAND	PALKDELLAE	ANVMQQLDNP	YIVRMIGICE	AESWMLVMEM	
AELGPLNKYL	QQNRHVKDKN	IIELVHQVSM	GMKYLEESNF	VHRDLAARNV	
LLVTQHYAKI	SDFGLSKALR	ADENYYKAQT	HGKWPVKWYA	PECINYYKFS	
SKSDVWSFGV	LMWEAFSYGQ	KPYRGMKGSE	VTAMLEKGER	MGCPAGCPRE	
MYDLMNLCWT	YDVENRPGFA	AVELRLRNYY	YDVVN		

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.