

RPG077Hu01 10µg

Recombinant Purinergic Receptor P2X, Ligand Gated Ion Channel 7 (P2RX7)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Ser47~Val334

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.6

Predicted Molecular Mass: 36.8kDa

Accurate Molecular Mass: 37kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[SEQUENCE]

				SDKL
YQRKEPVISS	VHTKVKGIAE	VKEEIVENGV	KKLVHSVFDT	ADYTFPLQGN
SFFVMTNFLK	TEGQEQRLCP	EYPTRRTLCS	SDRGCKKGWM	DPQSKGIQTG
RCVVYEGNQK	TCEVSAWCPI	EAVEEAPRPA	LLNSAENFTV	LIKNNIDFPG
HNYTTRNILP	GLNITCTFHK	TQNPQCPIFR	LGDIFRETGD	NFSDVAIQGG
IMGIEIYWDC	NLDRWFHHCR	PKYSFRRLDD	KTTNVSLYPG	YNFRYAKYYK
ENNVEKRTLI	KVFGIRFDIL	VFGTGGKFDI	IQLV	

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.