

RPJ828Hu01 100µg Recombinant Phosphoinositide-3-Kinase Class 3 (PIK3C3) Organism Species: Homo sapiens (Human) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression. Host: E. coli Residues: Tyr631~Trp885 Tags: N-terminal His-Tag Tissue Specificity: Testis, Uterus. Subcellular Location: Midbody. **Purity:** >98% Traits: Freeze-dried powder Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% Trehalose and Proclin300. Original Concentration: 200µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB. (May be suitable for use in other assays to be determined by the end user.) Predicted isoelectric point: 6.6 Predicted Molecular Mass: 32.9kDa Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

YPVIFKHGDD LRQDQLILQI ISLMDKLLRK ENLDLKLTPY KVLATSTKHG FMQFIQSVPV AEVLDTEGSI QNFFRKYAPS ENGPNGISAE VMDTYVKSCA GYCVITYILG VGDRHLDNLL LTKTGKLFHI DFGYILGRDP KPLPPPMKLN KEMVEGMGGT QSEQYQEFRK QCYTAFLHLR RYSNLILNLF SLMVDANIPD IALEPDKTVK KVQDKFRLDL SDEEAVHYMQ SLIDESVHAL FAAVVEQIHK FAQYW

[IDENTIFICATION]




Figure 1. SDS-PAGE