

RPA865Hu01 200µg

Recombinant Cluster Of Differentiation 247 (CD247)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Arg52~Arg164

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.5

Predicted Molecular Mass: 17.0kDa

Accurate Molecular Mass: 17kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

RVKFSRSAD APAYQQGQNQ LYNELNLGRR EEYDVLDKRR GRDPEMGGKP
QRRKNPQEGL YNELQKDKMA EAYSEIGMKG ERRRGKGDG LYQGLSTATK
DTYDALHMQA LPPR

[IDENTIFICATION]

ATGAGTGAAGTTCAGCAGGAGCGCAGCGCCCGGTACCAAGCAGGGCCAGAACCTCTATTAACGAGCTCAATCTAGGACGAAGAGAGGATACGATGTTTGGACAAAGAGACGTGGCCGGGACCCCTGAGATGGGGGAAAGCCGACAGAGGAAGAACCCCTCAGGAAGGCCGTGATCAJ
R V K F S R S A D A P A Y Q Q G Q N Q L Y N E L N L G R R E E Y D V L D K R R G R D P E M G G K P Q R R K N P Q E G L Y N




Figure. Gene Sequencing (Extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.