

RPB214Ra01 100µg

Recombinant Aspartate Aminotransferase (AST)

Organism Species: *Rattus norvegicus* (Rat)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Met1~Gln413

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.9

Predicted Molecular Mass: 47.6kDa

Accurate Molecular Mass: 44kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

MAPPSFFAQV PQAPPVLVFK LIADFRDDPD PRKVN LGVGA YRTDDSQPW
LPVVRKVEQK IANDHSLNHE YLPILGLAEF RSCASQLVLG DNSPALREN
VGGVQSLGGT GALRIGADFL GRWYNGTDNK NTPVYVSSPT WENHNGVFSA
AGFKDIRSYR YWDAEKRLD LQGFLNDLEN APEFSIFVLH ACAHNPTGTD
PTEEEWKQIA AVMKRRFLFP FFDSAYQGFA SGDLEKDAWA IRYFVSEGFE
LFCAQSFSKN FGLYNERVGN LTVVGKEHDS VLRVLSQMEK IVRITWSNPP
AQGARIVATT LSNPELFKEW KGNVKTADR ILTMRSELRA RLEALKTPGT
WSHITEQIGM FSFTGLNPKQ VEYLVNEKHI YLMPSGRINM CGLTTKNLDY
VATSINEAVT KFQ

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.