

RPD317Ra01 1mg

Recombinant Glycogen Synthase Kinase 3 Beta (GSK3b)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



## [ PROPERTIES ]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Lys122~Thr375

Tags: N-terminal His Tag

**Subcellular Location:** Membrane, Nucleus, Cytoplasm

**Purity:** > 97%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

0.01% SKL, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.8

Predicted Molecular Mass: 32.6kDa

Accurate Molecular Mass: 33kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [SEQUENCE]

		KKDEVYLNL	VLDYVPETVY	RVARHYSRAK
QTLPVIYVKL	YMYQLFRSLA	YIHSFGICHR	DIKPQNLLLD	PDTAVLKLCD
FGSAKQLVRG	EPNVSYICSR	YYRAPELIFG	ATDYTSSIDV	WSAGCVLAEL
LLGQPIFPGD	SGVDQLVEII	KVLGTPTREQ	IREMNPNYTE	FKFPQIKAHP
WTKVFRPRTP	PEAIALCSRL	LEYTPTARLT	PLEACAHSFF	DELRDPNVKL
PNGRDTPALF	NFTTQELSSN	PPLAT		

### [ IDENTIFICATION ]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.