

SAA544Cp08**FITC-Linked Rabbit Anti-Goat IgG Antibody****Organism Species: Capra hircus; Caprine (Goat)*****Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

[PROPERTIES]**Source:** Antibody labeling**Host:** Rabbit**Purification:** Antigen-specific Affinity Chromatography.**Label:** FITC**Traits:** Liquid**Concentration:** 200µg/mL**UOM:** 100µg**Applications:** WB; IF; IP; IHC-P; IHC-F; ELISA.**[IMMUNOGEN]****Immunogen:** IgG, Caprine**[APPLICATIONS]**

Western blotting: 1:2000-10000

Immunoprecipitation: 1:2000-10000

Immunohistochemistry in formalin fixed frozen section: 1:200-1000

Immunohistochemistry in paraffin section: 1:200-1000

Enzyme-linked Immunosorbent Assay: 1:4000-15000

Optimal working dilutions must be determined by end user.

[FORMULATION]**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**IDENTIFICATION**]




Figure 1: UV-spectrum

Sample: FITC-Linked Rabbit Anti-Goat IgG Ab (Catalog: SAA544Cp08)

FITC has a spectral characteristic at 495nm, which illustrate the success of FITC conjugation of antibody.