



Anti N-syndecan (syndecan-3)

BACKGROUND

Syndecan is the family name of transmembrane proteoglycans that carry predominantly heparan sulfate side chains. This proteoglycan family consists of four members. N-syndecan (syndecan-3) is the principal member expressed during early postnatal development in both central and peripheral nervous systems. N-syndecan binds various heparin-binding growth factors such as FGFs via the heparan sulfate moiety, and communicates with the cytoskeleton via the cytoplasmic domain of the core protein. N-syndecan plays a pivotal role in formation of the neural network through these molecular interactions. This antibody recognizes effectively the core protein of N-syndecan.

Product type	Primary antibody
Immunogen	A recombinant protein representing the entire core protein of rat N-syndecan
Raised in	-
Myeloma	-
Clone number	-
Isotype	-
Host	Rabbit
Source	Whole Serum (Liquid)
Purification	-
Buffer	PBS containing 0.1% NaN ₃ as a preservative
Concentration	-
Volume	100 ul
Label	Unlabeled
Specificity	N-syndecan core protein
Cross reactivity	Rat, Mouse
Storage	Shipped at 4°C. Upon arrival aliquot and store at -20°C or below. Aliquot to avoid cycles of freeze/thaw.
Other	Data Link : UniProtKB/Swiss-Prot P33671

Application notes	• Western blotting: 1/10,000
Recommended dilutions	• Immunohistochemistry: 1/10,000 (Paraffin) (Fig.2 / Ref.2)
	Other applications have not been tested. Optimal dilutions/concentrations should be determined by the end user.

References	1) Watanabe, E., Matsui, F., Keino, H., Ono, K., Kushima, Y., Noda, H., & Oohira, A., A membrane-bound heparan sulfate proteoglycan that is transiently expressed on growing axons in the rat brain, (1996) J. Neurosci. Res., 44, 84-96. 2) Toba, Y., Horie, M., Sango, K., Takashiki, A., Matsui, F., Oohira, A., & Kawano, H., Expression and immunohistochemical localization of heparan sulfate proteoglycan N-syndecan in the migratory pathway from the rat olfactory placode, (2002) Eur. J. Neurosci., 15, 1-13
-------------------	---

ANTIBODY CHARACTERIZATION




Fig.1 Characterization of the polyclonal anti-N-syndecan antiserum

Immunoblotting using anti-Nsyndecan of a partially purified preparation of membrane-bound proteoglycan (lane 1), and that digested with chondroitinase ABC (lane 2), and heparitinase I (lane 3). After the heparitinase I treatment, the immunoreactive band is sifted to an apparent molecular weight of approximately 140 kDa, identical to that of the core glycoprotein of rat N-syndecan.




Fig.2 Comparative localization of immunoreactivity for N-syndecan in the developing rat (16days) vomeronasal system.

Double immunofluorescence labelling with anti-NCAM (green) and anti-N-syndecan (red). N-syndecan immunoreactivity surrounds NCAM-immunoreactive vomeronasal axons and their associated cell clusters (asterisk).(Ref.2)

RELATED PRODUCTS:

Product Name	Maker	Cat#
Anti Chondroitin Sulfate A (2H6) Monoclonal Antibody	CAC	NU-07-001
Anti Neurocan (1G2) Monoclonal Antibody	CAC	NU-07-002
Anti Neuroglycan C (C1) Monoclonal Antibody	CAC	NU-07-003
Anti N-syndecan Polyclonal Antibody	CAC	NU-07-004
Anti Neurocan peptides Polyclonal Antibody	CAC	NU-07-005

For research use only. Not for clinical diagnosis.



COSMO BIO Co., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: <http://www.cosmobio.co.jp> e-mail: export@cosmobio.co.jp

[Outside Japan] Phone : +81-3-5632-9617 [国内連絡先] Phone : +81-3-5632-9610

FAX : +81-3-5632-9618 FAX : +81-3-5632-9619