



MONOCLONAL ANTIBODY

For research use only. Not for clinical diagnosis.

Catalog No. 2ZHIGG2

Anti Human IgG

Product type	Primary antibodies
Host	Mouse
Source	Ascites
Form	Hybridoma cell line (Hyb-IgG-19) was grown in ascitic fluid BALB/c Mouse Liquid Ammonium sulfate precipitation Physiological salt solution containing 0.1% Sodium Azide
Volume	0.5 mL
Concentration	0.5 mg/mL
Specificity	This antibody specifically reacts with Human IgG
Antigen	Purified Human IgG antigen
Clone	Hyb-IgG-19 The splenic lymphocytes from BALB/c mouse immunized with purified Human IgG were fused to mouse myeloma NS-1 cells and the hybrid cells (Hyb-IgG-19) were screened.
Isotype	IgG ₁

Application notes ELISA
Recommended use

Recommended dilutions

Optimal dilutions/concentrations should be determined by the end user.

Staining Pattern

Cross reactivity

Storage Maintain at 2-8 °C (-20 °C for long term storage) in undiluted aliquots.
Avoid repeated freeze/thaw cycles.

References 1) Takai E, et al: J.Immunol.Methods 95: 23-30,1986.





モノクローナル抗体

For research use only. Not for clinical diagnosis.

Catalog No. 2ZHIGG2

抗ヒト IgG

Product type	一次抗体
Host	マウス
Source	BALB/c マウス腹水 ハイブリドーマ (Hyb-IgG-19) を BALB/c マウスに接種して得られた腹水
Form	液状 硫酸アンモニウム塩析精製 0.1%アジ化ナトリウムを含む生理的食塩水溶液
Volume	0.5 mL
Concentration	0.5 mg/mL
Specificity	ヒト IgG に特異的に反応
Antigen	精製ヒト IgG 抗原
Clone	Hyb-IgG-19 精製ヒト IgG を免疫した BALB/c マウスの脾臓細胞とマウスミエローマ細胞 (NS-1) を融合して得られたハイブリドーマ
Isotype	IgG ₁

Application notes

ELISA

Recommended use

Recommended dilutions

Optimal dilutions/concentrations should be determined by the end user.

Staining Pattern

Cross reactivity

Storage

2~8°C 3ヶ月間

-20°C 3年間

(凍結融解の繰り返しは避けてください)

References

1) Takai E, et al: J.Immunol.Methods 95: 23-30,1986.



COSMO BIO Co., LTD.

Inspiration for Life Science

TOYO 2CHOME, KOTO-KU, TOKYO, 135-0016, JAPAN

URL: <http://www.cosmobio.co.jp>

e-mail: export@cosmobio.co.jp

[Outside Japan] Phone : +81-3-5632-9617

[国内連絡先] Phone : +81-3-5632-9610

FAX : +81-3-5632-9618

FAX : +81-3-5632-9619