



## Anti PACAP 27 (Human) Serum Cat. No. YII-Y050-EX Lot No. 080271013

**Description:** This antiserum was raised in a rabbit by immunization with a carrier free synthetic PACAP 27 (hu-man) peptide. The product vial contains 50 µL of the titled antiserum obtained by lyophilizing its 0.001 M phos- phate buffer (pH 7.0, 0.5mL) solution . It can be used for immunoassay, immunohistochemistry or any other immunoreaction with PACAP27 (human).

Immunogen: Synthetic PACAP 27 (human) peptide, carrier free Host: Rabbit

Amino Acid Sequence of PACAP27 (human)<sup>1)</sup>: HSDGIFTDSY SRYRKQMAVK KYLAAVL-NH2

Product Form: Lyophilized unpurified serum Size: 50 μL

**Reconstitution:** Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN3 0.1%).

**Storage:** The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon recon- stitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing- thawing should be avoided.

**Suggested Working Dilution Range:** 1:5,000-40,000 (final dilution ~1:280,000) for radioimmunoassay;

1: 1,000-4,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

**Specificity** (based on radioimmunoassay): PACAP27 (human) 100%, PACAP38 (human) 0%, VIP (porcine) 0%, PHI (rat) 0%, PHI (human) 0%, secretin (human) 0%, helodermin 0%

Positive Control (immunohistochemistry): Rat colon

Species Tested: Rat

## REFERENCES:

1) A. Miyata, A. Arimura et al., Isolation of a neuropeptide corresponding to the N-terminal 27 residues of the pituitary adenylate cyclase activating polypeptide with 38 residues (PACAP38). Biochemistry Biophysics Research Communications 170:643-648, 1990

## FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM

