



## Anti Pancreastatin (33-51) (Rat) Serum Cat. No. YII-Y090-EX Lot No. 087171031

**Description:** This antiserum was raised in a rabbit by immunization with a porcine thyroglobulin (pTG) conjugate of synthetic pancreastatin (33-51) (rat) peptide. The product vial contains 50 µL of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immuno- histochemistry or any other immunoreaction with pancreastatin (rat).

Immunogen: Synthetic pancreastatin (33-51) (rat)-pTG conjugate Host: Rabbit

Amino Acid Sequence of pancreastatin (33-51)(rat)<sup>1)</sup>:
DDGQSESQAV NGKTGASEAV PSEGKGELEH SQQEEDGEEAMAGPPQGLFP G

Product Form: Lyophilized unpurified serum Size: 50 µL

**Reconstitution:** Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN3 0.1%).

**Storage:** The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon recon- stitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing- thawing should be avoided.

Suggested Working Dilution Range: 1:10,000-200,000 (final dilution ~1:1,400,000) for radioimmunoassay;

1: 1,000- 4,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

**Specificity** (based on radioimmunoassay): pancreastatin (33-51) (rat) 100%, Pancreastatin (rat) 100%, chromo-granin A (359-389) (rat) 0%<sup>2)</sup>

Positive Control (immunohistochemistry): Rat pituitary and adrenal gland

Species Tested: Rat

## REFERENCES:

- 1) A. Iacangelo, H. Okayama and L.E. Eiden, Primary structure of rat chromogranin A and distribution of its mRNA. FEBS Letters 227:115-221, 1988
- 2) S. Nagasawa, N. Yanaihara et al., Distribution of chromogranin A-like peptides in the rat. Biomedical Research 16: 83-90, 1995

## FOR RESEARCH LABORATORY USE ONLY

DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM



TOYO EKIMAE BLDG, 2-20,TOYO 2CHOME,KOTO-KU,TOKYO 135-0016,JAPAN TEL: +81-3-5632-9617 FAX: +81-3-5632-9618 e-mail: export@cosmobio.co.jp www.cosmobio.co.jp