Elabscience®

CEBPA Polyclonal Antibody

catalog number: E-AB-64040

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human;Mouse;Rat |
| Immunogen | Recombinant fusion protein of human CEBPA (NP_004355.2). |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| WB | 1:500-1:2000 |
| IHC | 1:50-1:200 |

Data





Western blot analysis of extracts of Mouse lung using CEBPA Polyclonal Antibody at dilution of 1:1000.

Immunohistochemistry of paraffin-embedded Rat liver using CEBPA Polyclonal Antibody at dilution of 1:100 (40x lens).

Observed-MW:41 kDa Calculated-MW:25 kDa/35 kDa/37 kDa/41 kDa





Immunohistochemistry of paraffin-embedded Rat brain using Immunohistochemistry of paraffin-embedded Rat kidney CEBPA Polyclonal Antibody at dilution of 1:100 (40x lens). using CEBPA Polyclonal Antibody at dilution of 1:100 (40x lens).

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the |
| | temperature recommended. |

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017

Elabscience®

This intronless gene encodes a transcription factor that contains a basic leucine zipper (bZIP) domain and recognizes the CCAAT motif in the promoters of target genes. The encoded protein functions in homodimers and also heterodimers with CCAAT/enhancer-binding proteins beta and gamma. Activity of this protein can modulate the expression of genes involved in cell cycle regulation as well as in body weight homeostasis. Mutation of this gene is associated with acute myeloid leukemia. The use of alternative in-frame non-AUG (GUG) and AUG start codons results in protein isoforms with different lengths. Differential translation initiation is mediated by an out-of-frame, upstream open reading frame which is located between the GUG and the first AUG start codons.

For Research Use Only