

ARID3A antibody

Product Information

| | |
|---------------|---|
| Catalog No.: | FNab00566 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.) |

Background

This gene encodes a member of the ARID (AT-rich interaction domain) family of DNA binding proteins. It was found by homology to the Drosophila dead ringer gene, which is important for normal embryogenesis. Other ARID family members have roles in embryonic patterning, cell lineage gene regulation, cell cycle control, transcriptional regulation, and possibly in chromatin structure modification.


Immunogen information

| | |
|--------------|--|
| Immunogen: | AT rich interactive domain 3A (BRIGHT-like) |
| Synonyms: | AT-rich interactive domain-containing protein 3A (ARID domain-containing protein 3A) B-cell regulator of IgH transcription (Bright) Dead ringer-like protein 1 E2F-binding protein 1 ARID3A DRIL1 DRIL3 DRX E2FBP1 |
| Observed MW: | 100 kDa |
| Uniprot ID : | Q99856 |


Application

| | |
|-----------------------|--|
| Reactivity: | Human, Mouse, Rat |
| Tested Application: | ELISA, WB, IHC, IP |
| Recommended dilution: | WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IP: 1:20 - 1:50 |


Image:



Immunohistochemistry of paraffin-embedded human oophoroma using FNab00566(ARID3A antibody) at dilution of 1:100



IP Result of anti-ARID3A (IP:FNab00566, 4ug; Detection:FNab00566 1:1000) with K-562 cells lysate 3200ug.



K-562 cells were subjected to SDS PAGE followed by western blot with FNab00566(ARID3A antibody) at dilution of 1:1000