

## DTX3L antibody

## **Product Information**

Catalog No.: FNab02556

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months(Avoid repeated freeze / thaw cycles.)

## **Background**

Ubiquitin ligase that mediates monoubiquitination of 'Lys-91' of histone H4(H4K91ub1), in response to DNA damage. Protects cells exposed to DNA-damaging agents. The exact role of H4K91ub1 in DNA damage response is still unclear but it may function as a licensing signal for additional histone H4 post-translational modifications such as H4 'Lys-20' methylation(H4K20me). Involved in the recruitment of 53BP1/TP53BP1 to sites of DNA damage by mediating H4K91ub1 formation. In concert with PARP9, plays a role in PARP1-dependent DNA damage repair. PARP1-dependent PARP9-DTX3L-mediated ubiquitination promotes the rapid and specific recruitment of 53BP1/TP53BP1, UIMC1/RAP80, and BRCA1 to DNA damage sites.

## **Immunogen information**

Immunogen: deltex 3-like(Drosophila)

Synonyms: E3 ubiquitin-protein ligase DTX3L|B-lymphoma- and BAL-associated

protein|Protein deltex-3-like|RING-type E3 ubiquitin transferase

DTX3L|Rhysin-2 (Rhysin2)|DTX3L|BBAP

Observed MW: 84 kDa Uniprot ID: Q8TDB6

**Application** 


Reactivity: Human



Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



MCF-7 cells were subjected to SDS PAGE followed by western blot with FNab02556(DTX3L antibody) at dilution of 1:600