

HDAC4 antibody

Product Information

Catalog No.:	FNab09984
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3.


Immunogen information

Immunogen:	histone deacetylase 4
Synonyms:	Histone deacetylase 4 (HD4) HDAC4 KIAA0288
Observed MW:	130 kDa
Uniprot ID :	P56524

Application

Reactivity:	Human, Mouse
Tested Application:	ELISA, WB
Recommended dilution: WB:	1:500 - 1:2000

Image:



HeLa cells were subjected to SDS PAGE followed by western blot with FNab09984(HDAC4 antibody) at dilution of 1:1000