

**User Guide****HR2-432 (pg 1)****Application**

*In situ* proteolysis and proteolytic screening of protein samples for crystallization and structure determination

**Features**

- 6 proteases
- Stable, optimized, freeze dried protease formulation
- Enhanced stability
- Proti-Ace Dilution Buffer
- Optimized protocol for *in situ* proteolysis or proteolytic screening

**Kit Contents**

Proti-Ace 2 Reagent 1\* (Qty 3):

- 1 mg/ml Proteinase K,  
10 mM TRIS hydrochloride pH 7.5,  
1 mM Calcium acetate hydrate

Proti-Ace 2 Reagent 2\* (Qty 3):

- 1 mg/ml Clostripain (Endoproteinase-Arg-C),  
10 mM HEPES pH 7.5,  
2 mM TCEP hydrochloride,  
2 mM L-Cysteine,  
1 mM Calcium chloride dihydrate

Proti-Ace 2 Reagent 3\* (Qty 3):

- 1 mg/ml Pepsin,  
1 mM Hydrochloric acid

Proti-Ace 2 Reagent 4\* (Qty 3):

- 1 mg/ml Thermolysin,  
10 mM Calcium acetate hydrate pH 8.0,  
50 mM Sodium chloride

Proti-Ace 2 Reagent 5\* (Qty 3):

- 1 mg/ml Bromelain,  
5 mM HEPES pH 7.5,  
2 mM L-Cysteine,  
1 mM Ethylenediaminetetraacetic acid disodium salt dihydrate

Proti-Ace 2 Reagent 6\* (Qty 3):

- 1 mg/ml Actinase,  
5 mM HEPES pH 7.5,  
10 mM Sodium chloride

Proti-Ace Dilution Buffer (Qty 6):

- 1.6 ml of 10 mM HEPES pH 7.5,  
500 mM Sodium chloride

\* When 100 µl of deionized water is added to each Proti-Ace 2 freeze dried reagent

**Discussion**

A proteolytic fragment or domain of a protein may crystallize more readily or form better diffracting crystals than the intact protein.<sup>1-8</sup> Proteases can be used to generate small, active fragments or domains of the target protein for crystallization.<sup>9</sup> The fragment or domain can be used directly for crystallization experiments. Or the proteolytic sample analyzed by gel electrophoresis and/or mass spectrometry for mass and sequence for subsequent cloning, expression, purification and crystallization. Using proteolysis to enhance sample crystallization, the current overall success rate for yielding a deposited crystal structure is currently better than 12%.<sup>3</sup>

**Instructions for Proteolytic Screening**

Proteolytic screening is a procedure involving limited proteolysis of the sample versus a portfolio of proteases, followed by denaturing gel electrophoresis (SDS-PAGE) and/or mass spectrometry (MS) to identify regions of a gene corresponding to the protease resistant domain/fragment for subsequent cloning, expression, purification and crystallization.

1. Select all six, a subset or a single protease from the Proti-Ace 2 kit for Proteolytic Screening.
2. Add 100 µl of deionized water to each of the selected Proti-Ace 2 enzymes to create a 1 mg/ml Protease Stock solution.
3. Into empty micro centrifuge tubes, create a 1:100 dilution (0.01 mg/ml) of each 1 mg/ml Protease Stock from Step 2 by adding 5 µl of the 1 mg/ml Protease Stock plus 495 µl of Proti-Ace Dilution Buffer (10 mM HEPES pH 7.5, 500 mM Sodium chloride).
4. Pipette 10 µl of the 1:100 protease stock into 10 µl aliquots of protein (10 mg/ml) for each protease to be screened.
5. Incubate at 37°C for 60 minutes.
6. Stop the reaction by adding SDS-PAGE sample buffer for SDS-PAGE analysis or a final concentration of 10% v/v trichloroacetic acid for MS analysis. Refer to your SDS-PAGE and MS protocols for the appropriate volume and concentration of SDS-PAGE or MS sample buffer for quenching.
7. Analyze the digests by SDS-PAGE and/or MS. Identify the small active fragment (SAF) or protease resistant domains.<sup>9</sup> Clone the corresponding region of the gene. Express, purify and crystallize this gene product. Alternatively, scale up the proteolysis and purify the digest to produce a pure homogeneous sample of the SAF or domain for crystallization.

In the event of insufficient digestion, repeat steps 1-3 using a higher protease concentration such as 1:10 dilution of each Protease Stock (5 µl of 1 mg/ml Protease Stock plus 45 µl of Proti-Ace Dilution Buffer). Also consider longer incubation times, up to 24 hours.

In the event of over digestion, repeat steps 1-3 using a lower protease concentration such as 1:1,000 dilution of each Protease Stock (10 µl of the 1:100 Protease Stock plus 90 µl of Proti-Ace Dilution Buffer). Also consider shorter incubation times and/or lower incubation temperature (4 to 25°C).

## Instructions for *In Situ* Proteolysis

*In situ* proteolysis is a procedure where trace amounts of protease are included with the sample to be crystallized and mixed with crystallization reagents for screening or optimization experiments.<sup>2-4</sup>

1. Select the desired protease(s) from the Proti-Ace 2 kit to be used for *in situ* proteolysis.
2. Add 100 µl of deionized water to each of the selected Proti-Ace 2 enzymes to create a 1 mg/ml Protease Stock solution.
3. Add the protease to the protein crystallization sample. Add 10 µl of the 1 mg/ml Protease Stock solution to 90 µl of 10 mg/ml protein to create a 1:100 w/w dilution.
4. Set the crystallization experiment using the protease:sample mixture.

## Optimization of *In Situ* Proteolysis for Crystallization

- a. Vary the protease:sample ratio. Typical protease:sample ratios are 1:100, 1:1,000 and 1:10,000.
- b. Alter the incubation time. Typical incubation times are between 0 and 24 hours.
- c. Alter the incubation temperature. Typical incubation temperatures are between 4 and 37°C.
- d. For protein concentrations other than 10 mg/ml one can either use the preferred sample concentration with the protease:sample dilutions described in steps 1-4 or one can dilute the Proti-Ace 2 enzymes to a perfect 1:100 and/or 1:1,000 ratio based on the actual protein concentration. For example, if the protein concentration is 20 mg/ml one can add 50 µl of deionized water in step 2 to create a 2 mg/ml Protease Stock solution and then proceed with steps 3 and 4 to screen 1:100 protease:sample.

## Storage of the Proti-Ace 2 Kit

The unique freeze dried formulation of the Proti-Ace 2 kit offers a much improved protease stability compared to liquid protease formulations. Recommended storage: Room temperature up to 30 days, 4°C up to 12 months, -20°C up to 24 months. Once the proteases are made into solution the recommended storage is: Room temperature to 4°C up to 24 hours, -20°C up to 12 months.

## References

1. Allan D'Arcy, personal communication, 1989-2009
2. In situ proteolysis for protein crystallization and structure determination. Dong, A et al. Nature Methods - 4, 1019 - 1021 (2007)
3. In Situ Proteolysis to Generate Crystals for Structure Determination: An Update. Amy Wernimont, Aled Edwards. PLoS ONE 4(4): e5094. doi:10.1371/journal.pone.0005094
4. The use of in situ proteolysis in the crystallization of murine CstF-77. Tong et al. Acta Cryst. (2007). F63, 135-138
5. A brief history of protein crystal growth. McPherson, A. Journal of Crystal Growth, vol. 110, issue 1-2, pp. 1-10, 1991

6. Preparation and analysis of protein crystals. McPherson, A. John Wiley & Sons, Inc. 1982. ISBN 089464355X

7. A crystallizable form of the Streptococcus gordonii surface antigen SspB C-domain obtained by limited proteolysis. Forsgren et al. Acta Cryst. (2009). F65, 712-714

8. Preliminary X-ray analysis of a human VH fragment at 1.8 angstrom resolution. Gaur, Kupper, Fischer & Hoffman. Acta Cryst. (2004). D60, 965-967

9. Replication Protein A Characterization and Crystallization of the DNA Binding Domain. Pfuetzner et al. The Journal of Biological Chemistry, Vol. 272, No. 1, Issue of January 3, pp. 430-434, 1997.

## Related Products

**HR2-432-01** Proti-Ace 2 Reagent 1: 1 mg/ml Proteinase K, 10 mM TRIS hydrochloride pH 7.5, 1 mM Calcium acetate hydrate\*

**HR2-432-02** Proti-Ace 2 Reagent 2: 1 mg/ml Clostripain (Endoproteinase-Arg-C), 10 mM HEPES pH 7.5, 2 mM TCEP hydrochloride, 2 mM L-Cysteine, 1 mM Calcium chloride dihydrate\*

**HR2-432-03** Proti-Ace 2 Reagent 3: 1 mg/ml Pepsin, 1 mM Hydrochloric acid\*

**HR2-432-04** Proti-Ace 2 Reagent 4: 1 mg/ml Thermolysin, 10 mM Calcium acetate hydrate pH 8.0, 50 mM Sodium chloride\*

**HR2-432-05** Proti-Ace 2 Reagent 5: 1 mg/ml Bromelain, 5 mM HEPES pH 7.5, 2 mM L-Cysteine, 1 mM Ethylenediaminetetraacetic acid disodium salt dihydrate\*

**HR2-432-06** Proti-Ace 2 Reagent 6: 1 mg/ml Actinase, 5 mM HEPES pH 7.5, 10 mM Sodium chloride\*

**HR2-429-07** Proti-Ace Dilution Buffer: (10 mM HEPES pH 7.5, 500 mM Sodium chloride), 1.6 ml

**\* When 100 µl deionized water is added to the supplied freeze dried Proti-Ace 2 reagent.**

## Technical Support

Please e-mail (tech@hrmail.com), fax (1-949-425-1611), or telephone (1-949-425-1321 option 2) your request to Hampton Research. Fax and e-mail Technical Support are available 24 hours a day. Telephone technical support is available 8:00 a.m. to 4:30 p.m. USA Pacific Standard Time.

Hampton Research  
34 Journey

Aliso Viejo, CA 92656-3317 U.S.A.

Tel: (949) 425-1321 • Fax: (949) 425-1611

Technical Support e-mail: tech@hrmail.com

Website: www.hamptonresearch.com