

Recombinant Human HLA DRB1-14 beta

Source

Species	Human
Accession Number	Q9GIY3
Gene Symbol	HLA-DRB1
Expressed Region	Gly30-Lys227

Synonyms	HLA DRB1-14 beta, Major Histocompatibility Complex, Class II, DR Beta 1, DW2.2/DR2.2, DRw10, HLA Class II Histocompatibility Antigen, DR-1 Beta Chain, MHC Class II HLA-DR-Beta Cell Surface Glycoprotein, MHC Class II HLA-DR Beta 1 Chain, Human Leucocyte Antigen DRB1, MHC Class II Antigen DRB1*10, MHC Class II Antigen DRB1*11, MHC Class II Antigen DRB1*12, MHC Class II Antigen DRB1*13, MHC Class II Antigen DRB1*14, MHC Class II Antigen DRB1*15, MHC Class II Antigen DRB1*16, MHC Class II HLA-DRw10-Beta, MHC Class II Antigen DRB1*1, MHC Class II Antigen DRB1*3, MHC Class II Antigen DRB1*4, MHC Class II Antigen DRB1*7, MHC Class II Antigen DRB1*8, MHC Class II Antigen DRB1*9, Lymphocyte Antigen DRB1, MHC Class II Antigen, Clone P2-Beta-3, HLA-DR1B, HLA-DRB2, HLA-DRB, DR-12, DR-13, DR-14, DR-16, DRw11, DRB1, DR-1, DR-4, DR-5, DR-7, DR-8, DR-9, DR12, DR13, DR14, DR16, DRw8, DR5, DR4, SS1, DR1, DR7, DR8, DR9
-----------------	---

Preparation

Expression System	Human Embryonic Kidney 293 Cells
Tag	N-terminal 6x histidine tag
Purification	Unpurified cell culture supernatant. HEK293 cells grown in serum-free medium were transfected with expression vector harboring target gene. The cell culture was harvested with centrifugation to remove cells. The cell culture supernatant containing mammalian cell protease inhibitor cocktail was aliquoted and stored at -80 °C immediately. The gene overexpression in culture supernatant was confirmed by Western blotting using anti-His tag antibody and/or target-specific antibodies and the culture supernatant derived from HEK293 cells transfected with the empty expression vector was used as a negative control.
Molecular Weight	Recombinant Human HLA DRB1-14 beta has a calculated molecular mass of 23 kDa. The actual molecular weight may increase slightly due to the potential post-modifications (PTMs).

Protein Specifications

Format	Pink liquid
Formulation	Culture supernatant of transfected HEK293 cells
Preservative	None
Recommended Applications	Western blotting control, antibody validation (i.e., hybridoma screening, antibody pair test), ELISA, EIA, dot blotting, immunoprecipitation (IP), protein array, protein functional assay, protein-protein interaction, post-translational modifications, etc.

Shipping

Ice packs

Storage/Stability

Upon arrival, the protein may be stored for 2 weeks at 4 °C. For long term storage, it is recommended to store at -20 °C or -80 °C in appropriate aliquots. Avoid repeated freeze-thaw cycles.

This product is furnished for LABORATORY RESEARCH USE ONLY.

Not for diagnostic or therapeutic use.