

Recombinant Human TMEM123, transfected HEK293 cell culture supernatant**Source**

Species	Human
Accession Number	Q8N131-1
Gene Symbol	TMEM123
Expressed Region	His27-Thr166
Synonyms	TMEM123, Transmembrane Protein 123, Keratinocytes-Associated Transmembrane Protein 3, Pro-Oncosis Receptor Inducing Membrane Injury, KCT-3, KCT3, Pro Oncosis Receptor Inducing Membrane Injury Gene, Keratinocytes Associated Transmembrane Protein 3, Pro Oncosis Receptor Inducing Membrane Injury, Serine/Threonine-Rich Receptor, PORIMIN, PORMIN

Preparation

Expression System	Human Embryonic Kidney 293 Cells
Tag	N-terminal histidine tag
Purification	Unpurified cell culture supernatant. HEK293 cells grown in serum-free medium were transfected with expression vector harboring target gene. The cell culture was harvested with centrifugation to remove cells. The cell culture supernatant containing mammalian cell protease inhibitor cocktail was aliquoted and stored at -80 °C immediately. The gene overexpression in culture supernatant was confirmed by Western blotting using anti-His tag antibody and/or target-specific antibodies and the culture supernatant derived from HEK293 cells transfected with the empty expression vector was used as a negative control.
Molecular Weight	Recombinant protein has a calculated molecular mass of 14. The actual molecular weight may increase differently due to the potential post-modifications (PTMs).

Protein Specifications

Format	Pink liquid
Formulation	Culture supernatant of transfected HEK293 cells
Preservative	None
Recommended Applications	Western blotting control, antibody validation (i.e., hybridoma screening, antibody pair test), ELISA, EIA, dot blotting, immunoprecipitation (IP), protein array, protein functional assay, protein-protein interaction, post-translational modifications, etc.

Shipping

Ice packs

Storage/Stability

Upon arrival, the protein may be stored for 2 weeks at 4 °C. For long term storage, it is recommended to store at -20 °C or -80 °C in appropriate aliquots. Avoid repeated freeze-thaw cycles.

This product is furnished for LABORATORY RESEARCH USE ONLY.

Not for diagnostic or therapeutic use.