

3607 Parkway Lane, Suite 200 Peachtree Corners, GA 30092 www.RayBiotech.com | info@raybiotech.com 770-729-2992 | 1-888-494-8555 ISO 13485:2016 cGMP

Recombinant Human IFN receptor2 Protein, Biotinylated

Catalog Number: 230-30088B

Data Sheet / Certificate of Analysis (CoA)

Last Revised: 11/27/2024

Source

Species Human

Accession Number Q9BUA0

Gene Symbols IFNAR2

Synonyms IFN receptor2

Expressed Region Asp30-Gly246

Preparation

Expression System HEK293 cells

Tag N-terminal his-tag. The primary amino groups (-NH2) at the N-terminus and the side

chains of lysine (K) residues were biotin-conjugated using the standard chemical

labeling method.

Purity >95%

Molecular Weight Recombinant human IFNAR2 protein has a calculated molecular mass of 25 kDa.

Due to the abundant glycosylation, it migrates as approximately 35-45 kDa protein

bands in SDS-PAGE under DTT, beta-mercaptoethanol reducing conditions.

Specifications

Format Lyophilized powder

Formulation Lyophilized from a 0.2 ?m filtered solution in PBS

Concentration Determined by Pierce BCA protein assay kit

Reconstitution Briefly spin the vial and bring the contents to the bottom prior to opening. It is

recommended to reconstitute at 0.5 - 1.0 mg/mL with sterile deionized water.


For Research Use Only Page 1 of 2



Recombinant Human IFN receptor2 Protein, Biotinylated

Catalog Number: 230-30088B

SDS-PAGE Image



Shipping

The product is shipped with ice packs. Upon arrival, immediately store it at the temperature recommended below.

Storage/Stability

- Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
- Upon arrival, the lyophilized protein may be stored for 2 weeks at 4°C.
- For long term storage, it is recommended to store desiccated below -20°C in a manual defrost freezer.
- Generally, the shelf life is up to 12 months from date of receipt at -20°C or -80°C under sterile conditions.
- Following reconstitution, the protein may be stored for 2 weeks under sterile conditions at -20°C. For long term storage, it is recommended to make appropriate aliquots at -20°C or -80°C.

For Research Use Only Page 2 of 2