

Recombinant Equine IL-1RA

Source

- **Species** Equus caballus (Horse)
- **Gene Symbols** IL1RA
- **Accession Number** O18999
- **Expressed Region** His26-Gln177
- **Synonyms** Interleukin-1 receptor antagonist protein, IL-1RN, IL-1ra, IRAP, IL1 inhibitor.

Preparation

- **Expression System** Human embryonic kidney 293 (HEK293) cells
- **Tag** C-terminal his-tag
- **Purification** His-tag affinity purification by immobilized metal ion affinity chromatography (IMAC)
- **Purity** >95%
- **Endotoxin Level** <0.5 EU per µg of the protein as determined by the LAL method
- **Purity determined** By SDS-PAGE under reducing conditions and visualized by Coomassie blue staining
- **Molecular Weight** The recombinant protein product has a calculated molecular mass of 17 kDa. Due to the abundant glycosylation, it migrates as multiple large protein bands (17-40 kDa) in SDS-PAGE under DTT, beta-mercaptoethanol reducing conditions. After deglycosylation under denature condition, the protein presented as one single 17 kDa band. See deglycosylation analysis in SDS-PAGE image below.

Protein Specifications

- **Format** Liquid
- **Formulation** Supplied as a 0.2 µm filtered solution in PBS (pH 7.4)
- **Concentration** Lot specific (see the label on the vial), determined by BCA protein assay
- **SDS-PAGE Image**


Figure 1. Deglycosylation of purified recombinant proteins. The same amount of purified proteins were untreated (Lane 2) or treated with protein deglycosylation enzymes under native (Lane 3) or reducing (Lane 4) conditions. Deglycosylation treatment resulted in a mobility shift of the protein to produce one major band at the expected size, thus indicating that the untreated recombinant protein (Lane 2) was glycosylated. Lane 1: protein standard ladder (kDa).

Lane 2: untreated protein under reducing conditions.

Lane 3: treated protein with deglycosylation enzymes under native conditions.

Lane 4: treated protein with deglycosylation enzymes under denature conditions.

Lane 5: deglycosylation mixture only without target proteins.



Shipping

The product is shipped with ice packs

Storage/Stability

Upon arrival, the liquid protein may be stored for 2 weeks at 4°C. For long term storage, it is recommended to store at -20°C or -80°C in appropriate aliquots. Avoid repeated freeze-thaw cycles.

This product is furnished for **LABORATORY RESEARCH USE ONLY**.
Not for diagnostic or therapeutic use.