



Synonym


SNCA, NACP, PARK1, alpha-Synuclein

Source

Human Alpha-Synuclein (E46K), Tag Free(ALN-H5117) is expressed from E. coli cells. It contains AA Met 1 - Ala 140 (Accession # [P37840-1](#) (E46K)).

Predicted N-terminus: Met 1

Molecular Characterization



This protein carries no "tag".

The protein has a calculated MW of 14.5 kDa. The protein migrates as 16 kDa under reducing (R) condition (SDS-PAGE).

Purity

>95% as determined by SDS-PAGE.

Formulation

Supplied as 0.2 μ m filtered solution in 50mM HEPES, 100mM NaCl, pH8.0 with trehalose as protectant.

Contact us for customized product form or formulation.

Shipping

This product is supplied and shipped with dry ice, please inquire the shipping cost.


Storage

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- The product MUST be stored at -70°C or lower upon receipt;
- -70°C for 3 months under sterile conditions.

SDS-PAGE



Human Alpha-Synuclein (E46K), Tag Free on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

Discounts, Gifts,
and more!




» www.acrobiosystems.com

Human Alpha-Synuclein (E46K) Protein, Tag Free

Catalog # ALN-H5117



BIOSYSTEMS
Acro
Surprise Inside!



Immobilized Human Alpha-Synuclein (E46K), Tag Free (Cat. No. ALN-H5117) at 1 µg/mL (100 µL/well) can bind Synuclein alpha/beta Monoclonal Antibody with a linear range of 0.1-4 ng/mL (QC tested).

Background

Alpha-synuclein is a neuronal protein that plays several roles in synaptic activity such as regulation of synaptic vesicle trafficking and subsequent neurotransmitter release. It acts also as a molecular chaperone in its multimeric membrane-bound state, assisting in the folding of synaptic fusion components called SNAREs (Soluble NSF Attachment Protein REceptors) at presynaptic plasma membrane in conjunction with cysteine string protein-alpha/DNAJC5. Abnormalities in alpha-synuclein are implicated in the pathogenesis of Parkinson's disease (PD). Alpha-synuclein is present in Lewy-bodies, the neuropathological hallmark of PD, and the protein and its aggregation have been widely linked to neurotoxic pathways that ultimately lead to neurodegeneration.

Discounts, Gifts,
and more!



» www.acrobiosystems.com

9/3/2025