



## Synonym

ANGPT2, AGPT2, ANG2, Angiopoietin-2

## Source

Mouse Angiopoietin-2, His Tag(AN2-M5245) is expressed from human 293 cells (HEK293). It contains AA Lys 275 - Phe 496 (Accession # [O35608-1](#)).

Predicted N-terminus: His

## Molecular Characterization

Poly-his

ANGPT2(Lys 275 - Phe 496)  
O35608-1

This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 27.1 kDa. The protein migrates as 30-34 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

## Endotoxin

Less than 1.0 EU per  $\mu$ g by the LAL method / rFC method.

## Purity

>90% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

## Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in 20 mM MOPS, 150 mM NaCl, 5 mM CHAPS, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## SDS-PAGE



Mouse Angiopoietin-2, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

## SEC-MALS



The purity of Mouse Angiopoietin-2, His Tag (Cat. No. AN2-M5245) is more than 95% and the molecular weight of this protein is around 29-44 kDa verified by SEC-MALS.


[Report](#)

## Bioactivity-ELISA

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)



Immobilized Mouse Angiopoietin-2, His Tag (Cat. No. AN2-M5245) at 2 µg/mL (100 µL/well) can bind Human TIE2, Fc Tag (Cat. No. TI2-H5255) with a linear range of 0.039-0.625 µg/mL (QC tested).

## Background

Angiopoietin-2 is also known as ANGPT2, AGPT2, ANG2, and is a secreted glycoprotein that plays a complex role in angiogenesis and inflammation. Ang2 is widely expressed during development, but it is restricted postnatally to highly angiogenic tissues such as the placenta, ovaries, and uterus. It is particularly abundant in vascular endothelial cells (EC) where it is stored in intracellular Weibel Palade bodies. Both Ang2 and the related Angiopoietin1 (Ang1) are ligands for the receptor tyrosine kinase Tie 2. Ang2 functions as a proangiogenic factor, although it can also induce EC death and vessel regression. Upon its release from quiescent EC, it regulates vascular remodeling by promoting EC survival, proliferation, and migration and destabilizing the interaction between EC and perivascular cells. Ang2 is required for postnatal vascular remodeling, and it cooperates with Ang1 during lymphatic vessel development. It mediates the upregulation of ICAM1 and VCAM1 on EC, which facilitates the adhesion of leukocytes during inflammation. Ang2 competitively inhibits Ang1-induced endothelial cell responses mediated by Tie2, and reduces vascular integrity. But the role of Ang2 is controversial since the opposite outcomes have been reported in other studies. Over-expression of Ang2 disrupts the vascular remodeling, induces endothelial cell apoptosis, and may play an important regulating role in tumor angiogenesis. Ang2 also promotes the neuronal differentiation and migration of subventricular zone progenitor cells.

Discounts, Gifts,  
and more!

