

**Synonym**

ANGPTL3, ANGPT5, ANG-5, Angiopoietin-5, FHBL2

Source

Human Angiopoietin-like 3, His Tag(AN3-H5226) is expressed from human 293 cells (HEK293). It contains AA Ser 17 - Pro 220 (Accession # [NP_055310](#)).

Predicted N-terminus: Ser 17

Molecular Characterization

ANGPTL3(Ser 17 - Pro 220)	Poly-his
NP_055310	

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 24.6 kDa. The protein migrates as 28-35 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE


Human Angiopoietin-like 3, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

Discounts, Gifts,
and more!



» www.acrobiosystems.com



Immobilized Human Angiopoietin-like 3, His Tag (Cat. No. AN3-H5226) at 2 µg/mL (100 µL/well) can bind Monoclonal Anti-Human ANGPTL3 Antibody, Human IgG1 with a linear range of 0.2-6 ng/mL (QC tested).

Background

Angiopoietin-like protein 3 (ANGPTL3) is also known as Angiopoietin-related protein 3, Angiopoietin-5 (ANGPT5 / ANG-5), is a member of the angiopoietin-like family of secreted factors. ANGPTL3 / ANGPT5 is predominantly expressed in the liver, and has the characteristic structure of angiopoietins, consisting of a signal peptide, N-terminal coiled-coil domain and the C-terminal fibrinogen (FBN)-like domain. The FBN-like domain in angiopoietin-like 3 protein was shown to bind alpha-5/beta-3 integrins, and this binding induced endothelial cell adhesion and migration. This protein may also play a role in the regulation of angiogenesis. Angptl3 also acts as dual inhibitor of lipoprotein lipase (LPL) and endothelial lipase (EL), and increases plasma triglyceride and HDL cholesterol in rodents. ANGPTL3 inhibit endothelial lipase to catalyze HDL-phospholipid and increase HDL-PL levels. Circulating PL-riched HDL particles have high cholesterol efflux abilities.

Discounts, Gifts,
and more!



» www.acrobiosystems.com