



Synonym

B7-H6, NCR3LG1, B7 Homolog 6

Source

Human B7-H6 Protein, His Tag (B76-H52H8) is expressed from human 293 cells (HEK293). It contains AA Asp 25 - Ser 262 (Accession # [Q68D85-1](#)).

Predicted N-terminus: Asp 25

Molecular Characterization

B7-H6(Asp 25 - Ser 262)
Q68D85-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 28.6 kDa. The protein migrates as 38-55 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human B7-H6 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS



The purity of Human B7-H6 Protein, His Tag (Cat. No. B76-H52H8) is more than 85% and the molecular weight of this protein is around 37-55 kDa verified by SEC-MALS.


[Report](#)

Bioactivity-ELISA

Discounts, Gifts,
 and more!



» www.acrobiosystems.com



Immobilized Human NKp30, Fc Tag (Cat. No. NC3-H5259) at 10 µg/mL (100 µL/well) can bind Human B7-H6 Protein, His Tag (Cat. No. B76-H52H8) with a linear range of 0.016-0.25 µg/mL (QC tested).

Background

The B7 family of genes is essential in the regulation of the adaptive immune system. One of which is the recently discovered B7H6. Humans and rats have a single B7H6 gene; however, many B7H6 genes were detected in a single large cluster in the Xenopus genome.

Chimeric antigen receptor (CAR) T-cell therapies have demonstrated durable and potentially curative therapeutic efficacy against B-cell leukemia in clinical trials. In this study, B7H6, a ligand for the NK cell activating receptor NKp30, was targeted to create a CAR that targets multiple tumor types. B7H6 is expressed on various primary human tumors, including leukemia, lymphoma and gastrointestinal stromal tumors, but it is not constitutively expressed on normal tissues.

Discounts, Gifts,
and more!



» www.acrobiosystems.com