

Synonym


TNFRSF17, CD269, BCM, BCMA

Source

Biotinylated Cynomolgus / Rhesus macaque BCMA, His,Avitag (BCA-C82E5) is expressed from human 293 cells (HEK293). It contains AA Met 1 - Ala 53 (Accession # [A0A2K5UD97-1](#)). In the region Met 1 - Ala 53, the AA sequence of Cynomolgus and Rhesus macaque BCMA are homologous.

Predicted N-terminus: Met 1

Molecular Characterization



This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 9.6 kDa. The protein migrates as 15-18 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Cynomolgus / Rhesus macaque BCMA, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA

SEC-MALS



The purity of Biotinylated Cynomolgus / Rhesus macaque BCMA, His,Avitag (Cat. No. BCA-C82E5) is more than 90% and the molecular weight of this protein is around 12-17 kDa verified by SEC-MALS.


[Report](#)

Discounts, Gifts,
and more!




Biotinylated Cynomolgus / Rhesus macaque BCMA / TNFRSF17 Protein, His,Avitag™ (MALS verified)

Catalog # BCA-C82E5



Immobilized Human APRIL, His,Flag Tag, active trimer (Cat. No. APL-H52D1) at 2 µg/mL (100 µL/well) can bind Biotinylated Cynomolgus / Rhesus macaque BCMA, His,Avitag (Cat. No. BCA-C82E5) with a linear range of 0.2-1 ng/mL (QC tested).



Immobilized Biotinylated Cynomolgus / Rhesus macaque BCMA, His,Avitag (Cat. No. BCA-C82E5) at 1 µg/mL (100 µL/well) on streptavidin precoated (0.5 µg/well) plate, can bind Human BAFF, Fc Tag, active trimer (Cat. No. BAF-H5261) with a linear range of 0.2-5 ng/mL (Routinely tested).

Background

Tumor necrosis factor receptor superfamily member 17 (TNFRSF17) is also known as B-cell maturation protein (BCMA), CD antigen CD269, which is a member of the TNF-receptor superfamily. TNFRSF17 contains one TNFR-Cys repeat. TNFRSF17 is expressed in mature B-cells, but not in T-cells or monocytes. TNFRSF17 is receptor for TNFSF13B/BLyS/BAFF and TNFSF13/APRIL. TNFRSF17 promotes B-cell survival and plays a role in the regulation of humoral immunity. TNFRSF17 can activate NF-kappa-B and JNK.

