



Synonym

Cadherin-10, CDH10, T2-cadherin

Source

Human Cadherin-10, His Tag(CA0-H52H5) is expressed from human 293 cells (HEK293). It contains AA Gly 55 - Ala 613 (Accession # [Q9Y6N8-1](#)).

Molecular Characterization

CDH10(Gly 55 - Ala 613) Q9Y6N8-1	Poly-his
-------------------------------------	----------

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 64.2 kDa. The protein migrates as 70-80 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human Cadherin-10, His Tag on SDS-PAGE under reducing (R) condition.

The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Background

Cadherins are a large family of cell –cell adhesion molecules involved in inter-cellular adhesion in a wide variety of cell types. In the nervous system, cadherins are known to be crucial to all stages of development, including the early separation of the neural tube from the ectoderm, the segregation of neurones and axons, and the formation of synapses. These molecules mediate calcium ion-dependent, homotypic primers designed using two highly conserved amino acid sequences in the

Discounts, Gifts, and more!



» www.acrobiosystems.com

Human Cadherin-10 / CDH10 Protein, His Tag

Catalog # CA0-H52H5



BIOSYSTEMS
Acro
Surprise Inside!

cytoplasmic domain. One of the cadherins identified was cadherin-10. Cadherin-10 was first partially cloned from human brain, but its mRNA has also been shown to be present in mouse thymus (designated T2-cadherin), mouse testis, and in the developing brain and eye of mouse, rat, and chick.

Discounts, Gifts,
and more!



» www.acrobiosystems.com