



Synonym

CA125,CA-125,CA125MUC-16,FLJ14303,MUC16,mucin 16,mucin-16

Source

Biotinylated Human CA125 (13810-14451), Fc,Avitag(CA5-H82F8) is expressed from human 293 cells (HEK293). It contains AA Pro 13810 - Pro 14451 (Accession # [Q8WXI7-1](#)).  
Predicted N-terminus: Pro 13810

Molecular Characterization

|                                          |                                 |     |
|------------------------------------------|---------------------------------|-----|
| CA125(Pro 13810 - Pro 14451)<br>Q8WXI7-1 | Fc(Pro 100 - Lys 330)<br>P01857 | Avi |
|------------------------------------------|---------------------------------|-----|

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™).  
The protein has a calculated MW of 100.0 kDa. The protein migrates as 120-160 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 0.1 EU per µg by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.  
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.  
*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.  
*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Human CA125 (13810-14451), Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA





Immobilized Human Mesothelin (296-580), Fc Tag (Cat. No. MSN-H526x) at 5 µg/mL (100 µL/well) can bind Biotinylated Human CA125 (13810-14451), Fc,Avitag (Cat. No. CA5-H82F8) with a linear range of 0.6-78 ng/mL (QC tested).

Background

The CA125, also known as the MUC16, is a mucin protein that may be found in type I transmembrane or secreted forms that are used monitor the progress of epithelial ovarian cancer therapy. The CA 125 molecule is almost certainly a glycoprotein with a predominance of O-linkages. It is heterogeneous with regard to both size and charge, most likely due to continuous deglycosylation of side chains during its life-span in bodily fluids. It exists as a very large complex (perhaps as much as 4 million daltons) under natural conditions.

