

**Synonym**

CA125,CA-125,CA125MUC-16,FLJ14303,MUC16,mucin 16,mucin-16

Source

Rhesus macaque CA125 (13647-14291), His Tag(CA5-R52H6) is expressed from human 293 cells (HEK293). It contains AA Pro 13647 - Pro 14291 (Accession # [XP_015296314.1](#)).

Predicted N-terminus: Pro 13647

Molecular Characterization

CA125(Pro 13647 - Pro 14291)	Poly-his
XP_015296314.1	

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 73.9 kDa. The protein migrates as 95-140 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

EndotoxinLess than 0.1 EU per μ g by the LAL method / rFC method.**Purity**

>95% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

FormulationLyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Rhesus macaque CA125 (13647-14291), His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS

The purity of Rhesus macaque CA125 (13647-14291), His Tag (Cat. No. CA5-R52H6) is more than 95% and the molecular weight of this protein is around 90-110 kDa verified by SEC-MALS.


[Report](#)

Bioactivity-ELISA

Discounts, Gifts,
and more!



» www.acrobiosystems.com



Immobilized Rhesus macaque CA125 (13647-14291), His Tag (Cat. No. CA5-R52H6) at 5 µg/mL (100 µL/well) can bind Biotinylated Human Mesothelin, Fc,Avitag (Cat. No. MSN-H82F6) with a linear range of 0.002-0.313 µg/mL (QC tested).

Background

The CA125, also known as the MUC16, is a mucin protein that may be found in type I transmembrane or secreted forms that are used monitor the progress of epithelial ovarian cancer therapy. The CA 125 molecule is almost certainly a glycoprotein with a predominance of O-linkages. It is heterogeneous with regard to both size and charge, most likely due to continuous deglycosylation of side chains during its life-span in bodily fluids. It exists as a very large complex (perhaps as much as 4 million daltons) under natural conditions.

Discounts, Gifts,
and more!



» www.acrobiosystems.com