

**Synonym**

Cadherin-17,CDH17,HPT-1,LI-cadherin

**Source**

Biotinylated Mouse Cadherin-17 Protein, Fc,Avitag(CA7-M82F3) is expressed from human 293 cells (HEK293). It contains AA Phe 26 - Met 786 (Accession # [Q9R100-1](#)).

Predicted N-terminus: Phe 26

**Molecular Characterization**

Cadherin-17(Phe 26 - Met 786) Q9R100-1	Fc(Pro 100 - Lys 330) P01857	Avi
---	---------------------------------	-----

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 112.4 kDa. The protein migrates as 130-150 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

**Labeling**

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

**Protein Ratio**

Passed as determined by the HABA assay / binding ELISA.

**Purity**

&gt;90% as determined by SDS-PAGE.

**Formulation**

Lyophilized from 0.22 µm filtered solution in 20 mM Tris, 150 mM NaCl, pH 8.0 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*


**Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

**SDS-PAGE**

Biotinylated Mouse Cadherin-17 Protein, Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

**SEC-MALS**

The purity of Biotinylated Mouse Cadherin-17 Protein, Fc,Avitag (Cat. No. CA7-M82F3) is more than 85% and the molecular weight of this protein is around 270-310 kDa verified by SEC-MALS.


[Report](#)

**Bioactivity-ELISA**

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)



Immobilized Biotinylated Mouse Cadherin-17 Protein, Fc,Avitag (Cat. No. CA7-M82F3) at 1 µg/mL (100 µL/well) on streptavidin precoated (0.5 µg/well) plate. can bind Anti-Cadherin 17/CDH17 Antibody with a linear range of 0.002-0.03 % (QC tested).

## Background

Cadherin-17, also known as liver-intestine (LI) Cadherin, belongs to the cadherin family of calcium-dependent cell adhesion molecules. In vivo studies showed CDH17 knockout resulted in apoptotic PC tumor death through activating caspase-3 activity. Taken together, CDH17 functions as an oncogenic molecule critical to PC growth by regulating tumor apoptosis signaling pathways and CDH17 could be targeted to develop an anti-PC therapeutic approach.

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)