



Synonym

MDC,ABCD-1,SCYA22,STCP-1,DC,B-CK,A-152E5.1

Source

Human CCL22, Fc Tag(CC2-H5257) is expressed from human 293 cells (HEK293). It contains AA Gly 25 - Gln 93 (Accession # [O00626-1](#)).

Predicted N-terminus: Gly 25

Molecular Characterization

CCL22(Gly 25 - Gln 93)

O00626-1

Fc(Pro 100 - Lys 330)

P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 34.5 kDa. The protein migrates as 35-40 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, 0.2 M Arginine, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human CCL22, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Background

This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, dendritic cells, natural killer cells and for chronically activated T lymphocytes. It also displays a mild activity for primary activated T lymphocytes and has no chemoattractant activity for neutrophils, eosinophils and resting T lymphocytes. The product of this gene binds to

**Discounts, Gifts,
and more!**



» www.acrobiosystems.com

Human CCL22 / MDC Protein, Fc Tag

Catalog # CC2-H5257



BIOSYSTEMS
Acro
Surprise Inside!

chemokine receptor CCR4. This chemokine may play a role in the trafficking of activated T lymphocytes to inflammatory sites and other aspects of activated T lymphocyte physiology.

Discounts, Gifts,
and more!



» www.acrobiosystems.com