

**Synonym**

CD33, SIGLEC3, gp67

SourceCynomolgus Siglec-3, His Tag(CD3-C52H4) is expressed from human 293 cells (HEK293). It contains AA Met 16 - Leu 228 (Accession # [XP_045235686.1](#)).

Predicted N-terminus: Met 16

Molecular Characterization

Siglec-3(Met 16 - Leu 228)	Poly-his
XP_045235686.1	

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 25.7 kDa. The protein migrates as 38-43 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.**Endotoxin**Less than 1.0 EU per μ g by the LAL method / rFC method.**Purity**

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

FormulationLyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.


*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.***Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Cynomolgus Siglec-3, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS

The purity of Cynomolgus Siglec-3, His Tag (Cat. No. CD3-C52H4) is more than 90% and the molecular weight of this protein is around 40-60 kDa verified by SEC-MALS.

[Report](#)**Discounts, Gifts,
and more!**www.acrobiosystems.com

**Background**

Myeloid cell surface antigen CD33 is also known as SIGLEC3, Siglecs (sialic acid binding Iglike lectins) and GP67, is a single-pass type I membrane protein which belongs to the immunoglobulin superfamily and SIGLEC (sialic acid binding Ig-like lectin) family. Human CD33 / Siglec-3 cDNA encodes a 364 amino acid (aa) polypeptide with a hydrophobic signal peptide, an N-terminal Ig-like V-type domain, one Ig-like C2-type domains, a transmembrane region and a cytoplasmic tail. CD33 / Siglec-3 is usually considered myeloid-specific, but it can also be found on some lymphoid cells. In the immune response, CD33 / Siglec-3 may act as an inhibitory receptor upon ligand induced tyrosine phosphorylation by recruiting cytoplasmic phosphatase(s) via their SH2 domain(s) that block signal transduction through dephosphorylation of signaling molecules. CD33 / Siglec-3 induces apoptosis in acute myeloid leukemia.

**Discounts, Gifts,
and more!****» www.acrobiosystems.com**