


**Synonym**

CD5,LEU1

**Source**

Biotinylated Mouse CD5 Protein, His,Avitag(CD5-M82E3) is expressed from human 293 cells (HEK293). It contains AA Gln 24 - Asn 370 (Accession # [P13379](#)).

Predicted N-terminus: Gln 24

**Molecular Characterization**

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 41.4 kDa. The protein migrates as 53-63 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

The protein is designed as a dimer.

**Labeling**

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

**Protein Ratio**

Passed as determined by the HABA assay / binding ELISA.

**Purity**

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

**Formulation**

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*


**Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

**SDS-PAGE**

Biotinylated Mouse CD5 Protein, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

**SEC-MALS**

The purity of Biotinylated Mouse CD5 Protein, His,Avitag (Cat. No. CD5-M82E3) is more than 90% and the molecular weight of this protein is around 85-105 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)

**Background**

T-cell surface glycoprotein CD5 is also known as Lymphocyte antigen T1/Leu-1 and LEU1, which is phosphorylated on tyrosine residues by LYN, so CD5 can create binding sites for PTPN6/SHP-1. CD5 may act as a receptor in regulating T-cell proliferation. CD5 is expressed at various developmental and activation stages on human B cells. CD5 is a well established negative regulator of TCR and BCR signalling. CD5-positive cells may also prevent the emergence of autoimmunity by provision of cytokines such as IL-10. Development, selection and function of different B- and T-cell subsets or their preferential survival may be directly or indirectly dependent on different glycan structures associated with CD5 or CD5-like molecules.

**Discounts, Gifts,  
and more!****» [www.acrobiosystems.com](http://www.acrobiosystems.com)**