



Synonym

CD36, SCARB3, BDPLT10, CHDS7, FAT, GP3B, GP4, GPIV, PASIV, Platelet Glycoprotein 4, glycoprotein IV, gpIV, glycoprotein IIIb, gpIIIb

Source

Human CD36 Protein, Fc Tag(CD6-H5258) is expressed from human 293 cells (HEK293). It contains AA Gly 30 - Asn 439 (Accession # [P16671-1](#)).

Predicted N-terminus: Gly 30

Molecular Characterization

CD36(Gly 30 - Asn 439)	Fc(Pro 100 - Lys 330)
P16671-1	P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 73.1 kDa. The protein migrates as 95-120 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human CD36 Protein, Fc Tag on SDS-PAGE under reducing (R) condition.


The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

Discounts, Gifts,
and more!



» www.acrobiosystems.com



Immobilized Human CD36 Protein, Fc Tag (Cat. No. CD6-H5258) at 5 µg/mL (100 µL/well) can bind CD36 Monoclonal Antibody, Mouse/IgG1 with a linear range of 0.2-4 ng/mL (QC tested).

Background

CD36 (Cluster of Differentiation 36) is also known as platelet membrane glycoprotein IV (GPIV), fatty acid translocase (FAT), thrombospondin receptor, collagen receptor, and scavenger receptor class B, member 3 (SRB3), is a member of the class B scavenger receptor family of cell surface proteins. The human CD36 gene encodes a single chain 472 amino acid residue protein containing both an N- and a C-terminal cytoplasmic tail and an extracellular loop. CD36 is found on platelets, erythrocytes, monocytes, differentiated adipocytes, mammary epithelial cells, spleen cells and some skin microdermal endothelial cells. CD36 is a multiligand pattern recognition receptor that interacts with a large number of structurally dissimilar ligands, including long chain fatty acid (LCFA), advanced glycation end products (AGE), thrombospondin-1, oxidized low-density lipoproteins (oxLDLs), high density lipoprotein (HDL), phosphatidylserine, apoptotic cells, beta-amyloid fibrils (fA β), collagens I and IV, and Plasmodium falciparum infected erythrocytes. CD36 is required for the anti-angiogenic effects of thrombospondin1 In the corneal neovascularization assay. On binding a ligand the protein and ligand are internalized. This internalization is independent of macropinocytosis and occurs by an actin dependent mechanism requiring the activation Src-family kinases, JNK and Rho-family GTPases. CD36 ligands have also been shown to promote sterile inflammation through assembly of a Toll-like receptor 4 and 6 heterodimer.

Discounts, Gifts,
and more!



» www.acrobiosystems.com