

**Synonym**

CD117,SCFR,c-Kit,KIT

Source

Human CD117, His Tag(CD7-H52H4) is expressed from human 293 cells (HEK293). It contains AA Gln 26 - Thr 516 (Accession # [P10721-2](#)).

Predicted N-terminus: Gln 26

Molecular Characterization

CD117(Gln 26 - Thr 516)
P10721-2

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 57.1 kDa. The protein migrates as 80-95 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Human CD117, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS

The purity of Human CD117, His Tag (Cat. No. CD7-H52H4) is more than 90% and the molecular weight of this protein is around 75-100 kDa verified by SEC-MALS.


[Report](#)

Bioactivity-ELISA

Discounts, Gifts,
and more!




» www.acrobiosystems.com



Immobilized Human CD117, His Tag (Cat. No. CD7-H52H4) at 2 µg/mL (100 µL/well) can bind Biotinylated Human SCF, Avitag, His Tag (Cat. No. SCF-H82E1) with a linear range of 0.5-8 ng/mL (QC tested).

Bioactivity-SPR



Biotinylated Human SCF, Avitag, His Tag (Cat. No. SCF-H82E1) captured on Biotin CAP-Series S Sensor Chip can bind Human CD117, His Tag (Cat. No. CD7-H52H4) with an affinity constant of 0.236 µM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Background

Mast/stem cell growth factor receptor Kit (c-Kit), a member of the protein kinase superfamily, Tyr protein kinase family and CSF-1/PDGF receptor subfamily, is also known as piebald trait protein (PBT), p145 c-Kit, tyrosine-protein kinase Kit and CD117, which contains five Ig-like C2-type (immunoglobulin-like) domains and one protein kinase domain. CD117 acts as cell-surface receptor for the cytokine KITLG/SCF and plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and function, and in melanogenesis. In response to KITLG/SCF binding, CD117 can activate several signaling pathways. Furthermore, Activated KIT promotes phosphorylation of the protein phosphatases PTPN6/SHP-1 and PTPRU, and of the transcription factors STAT1, STAT3, STAT5A and STAT5B.

Discounts, Gifts,
and more!



» www.acrobiosystems.com