

**Synonym**

CD117,SCFR,c-Kit,KIT

Source

Rhesus macaque CD117, Fc Tag(CD7-R5259) is expressed from human 293 cells (HEK293). It contains AA Gln 26 - Thr 520 (Accession # [NP_001253024.1](#)).

Predicted N-terminus: Gln 26

Molecular Characterization

CD117(Gln 26 - Thr 520)	Fc(Pro 100 - Lys 330)
NP_001253024.1	P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 82 kDa. The protein migrates as 100-120 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

EndotoxinLess than 1.0 EU per μ g by the LAL method / rFC method.**Purity**

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Rhesus macaque CD117, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS

The purity of Rhesus macaque CD117, Fc Tag (Cat. No. CD7-R5259) is more than 85% and the molecular weight of this protein is around 188-228 kDa verified by SEC-MALS.



[Report](#)

Bioactivity-ELISA

Discounts, Gifts,
and more!



» www.acrobiosystems.com

**Rhesus macaque CD117, Fc Tag ELISA**
0.2 µg of Rhesus macaque CD117, Fc Tag per well**Rhesus macaque CD117, Fc Tag ELISA**
0.2 µg of ActiveMax Human SCF, His Tag per well

Immobilized Rhesus macaque CD117, Fc Tag (Cat. No. CD7-R5259) at 2 µg/mL (100 µL/well) can bind Human SCF protein, His Tag with a linear range of 0.2-2 ng/mL (QC tested).

Immobilized Human SCF protein, His Tag at 2 µg/mL (100 µL/well) can bind Rhesus macaque CD117, Fc Tag (Cat. No. CD7-R5259) with a linear range of 0.3-10 ng/mL (Routinely tested).

Background

Mast/stem cell growth factor receptor Kit (c-Kit), a member of the protein kinase superfamily, Tyr protein kinase family and CSF-1/PDGF receptor subfamily, is also known as piebald trait protein (PBT), p145 c-kit, tyrosine-protein kinase Kit and CD117, which contains five Ig-like C2-type (immunoglobulin-like) domains and one protein kinase domain. CD117 acts as cell-surface receptor for the cytokine KITLG/SCF and plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and function, and in melanogenesis. In response to KITLG/SCF binding, CD117 can activate several signaling pathways. Furthermore, Activated KIT promotes phosphorylation of the protein phosphatases PTPN6/SHP-1 and PTPRU, and of the transcription factors STAT1, STAT3, STAT5A and STAT5B.

Discounts, Gifts,
and more!



» www.acrobiosystems.com