



Synonym

CD28, Tp44

Source

Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag (CD8-H82F2) is expressed from human 293 cells (HEK293). It contains AA Asn 19 - Pro 152 (Accession # [AAH93698](#)). In the region Asn 19 - Pro 152, the AA sequence of Human, Cynomolgus and Rhesus macaque CD28 are homologous. Predicted N-terminus: Asn 19

Molecular Characterization

CD28(Asn 19 - Pro 152)	Fc(Pro 100 - Lys 330)	Avi
AAH93698	P01857	

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 43.6 kDa. The protein migrates as 55-70 kDa under reducing (R) condition, and 110-130 kDa under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>95% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS



The purity of Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag (Cat. No. CD8-H82F2) is more than 95% and the molecular weight of this protein is around 133-163 kDa verified by SEC-MALS.

[Report](#)

Bioactivity-ELISA


Discounts, Gifts,
and more!




» www.acrobiosystems.com



Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag ELISA
0.1 µg of Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag per well




Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag ELISA
0.5 µg of Anti-CD28 MAb, Mouse IgG1 per well




Immobilized Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag (Cat. No. CD8-H82F2) at 1 µg/mL (100 µL/well) on Streptavidin (Cat. No. STN-N5116) precoated (0.5 µg/well) plate, can bind Human B7-1, Mouse IgG2a Fc Tag, low endotoxin (Cat. No. B71-H52A4) with a linear range of 5-78 ng/mL (QC tested).

Immobilized Anti-CD28 MAb, Mouse IgG1 at 5 µg/mL (100 µL/well) can bind Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag (Cat. No. CD8-H82F2) with a linear range of 2.4-78 ng/mL (Routinely tested).

Bioactivity-BLI



Loaded Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag (Cat. No. CD8-H82F2) on SA Biosensor, can bind Human B7-1, Fc Tag (Cat. No. B71-H5259) with an affinity constant of 56.8 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Biotinylated Human / Cynomolgus / Rhesus macaque CD28, Fc,Avitag (Cat. No. CD8-H82F2) on SA Biosensor, can bind Human B7-1, Mouse IgG2a Fc Tag (Cat. No. B71-H52A4) with an affinity constant of 38.3 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Background

T-cell-specific surface glycoprotein CD28 is also known as TP44, is a single-pass type I membrane protein which contains one Ig-like V-type (immunoglobulin-like) domain. is one of the molecules expressed on T cells that provide co-stimulatory signals, which are required for T cell activation. CD28 is the receptor for CD80 (B7.1) and CD86 (B7.2). When activated by Toll-like receptor ligands, the CD80 expression is upregulated in antigen presenting cells (APCs). The CD86 expression on antigen presenting cells is constitutive. CD28 is the only B7 receptor constitutively expressed on naive T cells.

**Discounts, Gifts,
and more!**

