

**Synonym**

CD28, Tp44

Source

Mouse CD28, His Tag(CD8-M52H6) is expressed from human 293 cells (HEK293). It contains AA Asn 20 - Lys 149 (Accession # [P31041-1](#)).

Predicted N-terminus: Asn 20

Molecular Characterization

CD28(Asn 20 - Lys 149)
P31041-1 **Poly-his**

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 16.9 kDa. The protein migrates as 30-38 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Mouse CD28, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS

The purity of Mouse CD28, His Tag (Cat. No. CD8-M52H6) is more than 90% and the molecular weight of this protein is around 53-63 kDa verified by SEC-MALS.


[Report](#)

Bioactivity-ELISA

Discounts, Gifts,
 and more!



» www.acrobiosystems.com



Immobilized Mouse CD28, His Tag (Cat. No. CD8-M52H6) at 5 µg/mL (100 µL/well) can bind Mouse B7-1, Fc Tag (Cat. No. CD0-M5259) with a linear range of 0.02-0.156 µg/mL (QC tested).

Background

T-cell-specific surface glycoprotein CD28 is also known as TP44, is a single-pass type I membrane protein which contains one Ig-like V-type (immunoglobulin-like) domain. is one of the molecules expressed on T cells that provide co-stimulatory signals, which are required for T cell activation. CD28 is the receptor for CD80 (B7.1) and CD86 (B7.2). When activated by Toll-like receptor ligands, the CD80 expression is upregulated in antigen presenting cells (APCs). The CD86 expression on antigen presenting cells is constitutive. CD28 is the only B7 receptor constitutively expressed on naive T cells.

Discounts, Gifts,
and more!



» www.acrobiosystems.com