



Synonym

CD39,ENTPD1,NTPDase 1,Entpd1,Ecto-ATPDase 1,Ecto-ATPase 1

Source

Human CD39 (Q96A, N99A, E143A, R147E) Protein, His Tag(CD9-H52H5) is expressed from human 293 cells (HEK293). It contains AA Thr 38 - Val 478 (Accession # [P49961-1](#) (Q96A, N99A, E143A, R147E)).

Molecular Characterization

Q96A, N99A, E143A, R147E

**CD39 (Thr 38 - Val 478)
P49961-1**

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 52.1 kDa. The protein migrates as 60-70 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Supplied as 0.2 μ m filtered solution in 20 mM Tris,150 mM NaCl,pH8.0 with trehalose as protectant.

Contact us for customized product form or formulation.

Shipping

This product is supplied and shipped with dry ice, please inquire the shipping cost.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human CD39 (Q96A, N99A, E143A, R147E) Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS



The purity of Human CD39 (Q96A, N99A, E143A, R147E) Protein, His Tag (Cat. No. CD9-H52H5) is more than 90% and the molecular weight of this protein is around 65-85 kDa verified by SEC-MALS.

[Report](#)

Bioactivity-ELISA


Discounts, Gifts,
and more!



» www.acrobiosystems.com



Human CD39 / ENTPD1 (Q96A, N99A, E143A, R147E), His Tag ELISA
0.1 µg of Human CD39 / ENTPD1 (Q96A, N99A, E143A, R147E), His Tag per well



Immobilized Human CD39 / ENTPD1 (Q96A, N99A, E143A, R147E), His Tag (Cat. No. CD9-H52H5) at 1 µg/mL (100 µL/well) can bind X-IgG1 with a linear range of 0.5-3 ng/mL (Routinely tested).

Bioactivity

Measured by its ability to hydrolyze the 5'-phosphate group from the substrate adenosine-5'-triphosphate (ATP). The specific activity is > 8,000 pmol/min/µg (QC tested).

Background

CD39 is also known as Ectonucleoside triphosphate diphosphohydrolase 1, ENTPD1, NTPDase 1, Ecto-ATPDase 1, in the nervous system, could hydrolyze ATP and other nucleotides to regulate purinergic neurotransmission. Could also be implicated in the prevention of platelet aggregation by hydrolyzing platelet-activating ADP to AMP. Hydrolyzes ATP and ADP equally well. NTPDase-1 was originally described as CD39, a B lymphocyte cell surface marker, but it is also present on the surface of natural killer cells, T cells, and some endothelial cells. Regulatory T cells (Tregs) mediate immunosuppression through multiple, non-redundant, cell-contact dependent and independent mechanisms, a growing body of evidence suggests an important role for the CD39-CD73-adenosine pathway. CD39 ectonucleotidase is the rate-limiting enzyme of a cascade leading to the generation of suppressive adenosine that alters CD4 and CD8 T cell and natural killer cell antitumor activities.

Discounts, Gifts,
and more!



» www.acrobiosystems.com