

**Synonym**

FCGR

Source

Cynomolgus CD32b, His Tag(CDB-C5224) is expressed from human 293 cells (HEK293). It contains AA Ala 46 - Pro 224 (Accession # [Q8SPW3-1](#)).

Predicted N-terminus: Ala 46

Molecular Characterization

CD32b(Ala 46 - Pro 224)	Poly-his
Q8SPW3-1	

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 21.8 kDa. The protein migrates as 30-35 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 12 months under sterile conditions after reconstitution.

SDS-PAGE

Cynomolgus CD32b, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

SEC-MALS

The purity of Cynomolgus CD32b, His Tag (Cat. No. CDB-C5224) is more than 95% and the molecular weight of this protein is around 23-35 kDa verified by SEC-MALS.


[Report](#)

Bioactivity-SPR

Discounts, Gifts,
and more!



» www.acrobiosystems.com



Cynomolgus CD32b, His Tag (Cat. No. CDB-C5224) captured on CM5 chip via anti-His antibody can Herceptin® with an affinity constant of 11.9 μ M as determined in a SPR assay (Biacore T200) (QC tested).

Background

Receptors for the Fc region of IgG (Fc γ R) are members of the Ig superfamily that function in the activation or inhibition of immune responses. Three classes of human Fc γ Rs: RI (CD64), RII (CD32), and RIII (CD16), which generate multiple isoforms, are recognized.

There are three genes for human Fc γ RII /CD32 (A, B, and C) and one for mouse Fc γ RII B (CD32B). CD32 is a low affinity receptor for IgG. Low affinity immunoglobulin gamma Fc region receptor II-b (FCGR2B) is also known as CD32b, FCG2, IGFR2. CD32B is expressed on B cells and myeloid dendritic cells. Ligation of CD32B on B cells downregulates antibody production and may, in some circumstances, promote apoptosis. Co-ligation of CD32B on dendritic cells inhibits maturation and blocks cell activation. CD32B may also be a target for monoclonal antibody therapy for malignancies.

Discounts, Gifts,
and more!



» www.acrobiosystems.com