

**Synonym**

FCGR2B,C,CD32b,c,FcRII-b,c,Fc-gamma RII-b,c,Fc-gamma-RIIb,c,CD32,FCG2,IGFR2,CDw32

**Source**

Human CD32b/c Protein, Strep Tag(CDB-H5284) is expressed from human 293 cells (HEK293). It contains AA Ala 46 - Pro 217 (Accession # [P31994-1](#)).

Predicted N-terminus: Ala 46

**Molecular Characterization**

CD32b/c(Ala 46 - Pro 217)	Twin-Strep
P31994-1	

This protein carries a twin strep tag at the C-terminus.

The protein has a calculated MW of 22.7 kDa. The protein migrates as 30-34 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

**Endotoxin**

Less than 1.0 EU per  $\mu$ g by the LAL method / rFC method.

**Purity**

>95% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

**Formulation**

Lyophilized from 0.22  $\mu$ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*


**Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

**SDS-PAGE**

Human CD32b/c Protein, Strep Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

**SEC-MALS**

The purity of Human CD32b/c Protein, Strep Tag (Cat. No. CDB-H5284) is more than 95% and the molecular weight of this protein is around 26-38 kDa verified by SEC-MALS.


[Report](#)

**Bioactivity-SPR**


Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)



Human CD32b/c Protein, Strep Tag (Cat. No. CDB-H5284) immobilized on CM5 Chip can bind Rituximab with an affinity constant of 10.2  $\mu$ M as determined in a SPR assay (Biacore 8K) (QC tested).



Rituximab immobilized on CM5 Chip can bind Human CD32b/c Protein, Strep Tag (Cat. No. CDB-H5284) with an affinity constant of 5.18  $\mu$ M as determined in a SPR assay (Biacore 8K) (Routinely tested).

## Background

Receptors for the Fc region of IgG (Fc  $\gamma$  R) are members of the Ig superfamily that function in the activation or inhibition of immune responses. Three classes of human Fc  $\gamma$  Rs: RI (CD64), RII (CD32), and RIII (CD16), which generate multiple isoforms, are recognized.

There are three genes for human Fc $\gamma$  RII /CD32 (A, B, and C) and one for mouse Fc $\gamma$  RII B (CD32B). CD32 is a low affinity receptor for IgG. Low affinity immunoglobulin gamma Fc region receptor II-b (FCGR2B) is also known as CD32b, FCG2, IGFR2. CD32B is expressed on B cells and myeloid dendritic cells. Ligation of CD32B on B cells downregulates antibody production and may, in some circumstances, promote apoptosis. Co-ligation of CD32B on dendritic cells inhibits maturation and blocks cell activation. CD32B may also be a target for monoclonal antibody therapy for malignancies.

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)