


Synonym

Fc gamma RIIB,CD16b
(NA2),FCGR3B,CD16B,FCG3B,FCGR3,FCG3,IGFR3

Source

Human CD16b (NA2), SUMO,His Tag (CDB-H5294) is expressed from human 293 cells (HEK293). It contains AA Gly 17 - Ser 200 (Accession # [O75015-1](#)). The NA1 form of the CD16b differ with the NA2 form of CD16b in AA36, 65, 82, and 106. The NA1 form carries R36, N65, D82, and V106, while the NA2 form carries S36, S65, N82, and I106.
Predicted N-terminus: Gly

Molecular Characterization



This protein carries a SUMO tag at the N-terminus and a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 33.5 kDa. The protein migrates as 45-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.
>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.
This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 12 months under sterile conditions after reconstitution.

SDS-PAGE



Human CD16b (NA2), SUMO,His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.


SEC-MALS



The purity of Human CD16b (NA2), SUMO,His Tag (Cat. No. CDB-H5294) is more than 95% and the molecular weight of this protein is around 40-55 kDa verified by SEC-MALS.
[Report](#)

Bioactivity-BLI





Loaded Human CD16b (NA2), SUMO,His Tag (Cat. No. CDB-H5294) on HIS1K Biosensor, can bind Rituximab with an affinity constant of 4.8 μ M as determined in BLI assay (ForteBio Octet Red96e) (QC tested).

Background

CD16 is a low affinity Fc receptor, and has been identified as Fc receptors Fc γ RIIIa (CD16a) and Fc γ RIIIb (CD16b). These receptors bind to the Fc portion of IgG antibodies. CD16 encoded by two different highly homologous genes in a cell type-specific manner.CD16 is found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages.

CD16B is also kown as FCGR3B and FCG3B, is expressed specifically by polymorphonuclear leukocytes (neutrophils) and stimulated eosinophils. CD16B is the low affinity receptor for the Fc region of immunoglobulins gamma. FCGR3B binds complexed or aggregated IgG and also monomeric IgG. Contrary to III-A, FCG3B is not capable to mediate antibody-dependent cytotoxicity and phagocytosis. CD16B may serve as a trap for immune complexes in the peripheral circulation which does not activate neutrophils.

