



Synonym

CD70, CD27LG, TNFSF7, TNFSF7G, CD27L

Source

Human CD27 Ligand Protein, Fc Tag (CDL-H5266) is expressed from human 293 cells (HEK293). It contains AA Ser 52 - Pro 193 (Accession # [P32970-1](#)) trimer Design.

Predicted N-terminus: Pro

Molecular Characterization

This protein carries a human IgG1 Fc tag at the N-terminus.

The protein has a calculated MW of 74.1 kDa. The protein migrates as 130-150 kDa under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in Tris, Glycine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human CD27 Ligand Protein, Fc Tag on SDS-PAGE under non-reducing (NR) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

SEC-MALS



The purity of Human CD27 Ligand Protein, Fc Tag (Cat. No. CDL-H5266) is more than 85% and the molecular weight of this protein is around 160-190 kDa verified by SEC-MALS.


[Report](#)

Bioactivity-ELISA


Discounts, Gifts,
and more!



» www.acrobiosystems.com



Immobilized Human CD27, His Tag (Cat. No. CD7-H522b) at 2 µg/mL (100 µL/well) can bind Human CD27 Ligand Protein, Fc Tag (Cat. No. CDL-H5266) with a linear range of 0.2-25 ng/mL (QC tested).



Serial dilutions of Anti-CD27 Ligand (cusatuzumab) Antibody, Human IgG1 were added into Human CD27 Ligand Protein, Fc Tag (Cat. No. CDL-H5266): Biotinylated Human CD27, Fc,Avitag (Cat. No. TN7-H82F6) binding reactions. The half maximal inhibitory concentration (IC50) is 0.74034 µg/mL (Routinely tested).

Background

Cluster of Differentiation 70 (CD70) is also known as CD27 ligand (CD27L / CD27LG), TNFSF7, TNFSF7G, is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and is a surface antigen found on activated T-and B-lymphocytes and mature dendritic cells. Binding of CD70 to its receptor CD27 induces in priming, effector functions, differentiation and memory formation of T-cells, and thus is involved in the biological processes including T-cell activation, the proliferation of costimulated T-cells, as well as the generation of cytolytic T-cells. CD70 on T cells provides costimulatory signals that are required for T cell proliferation, clonal expansion and the promotion of effector T cell formation. CD70 on mouse B cell has been shown to inhibit terminal differentiation of activated B cells into plasma cells and enhances commitment to memory B cell responses. CD70 induces proliferation and IFN γ production, on NK cells.

Discounts, Gifts,
and more!

