

**Synonym**

CEACAM1,CD66a,BGP,BGP1,BGPI

Source

Human CEACAM-1, His Tag(CE1-H5220) is expressed from human 293 cells (HEK293). It contains AA Gln 35 - Gly 428 (Accession # [AAH14473](#)).

Predicted N-terminus: Gln 35

Molecular Characterization

CEACAM-1(Gln 35 - Gly 428)
AAH14473

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 44.1 kDa. The protein migrates as 57-100 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to different glycosylation.

Endotoxin

Less than 0.1 EU per μ g by the LAL method / rFC method.

Purity

>92% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Human CEACAM-1, His Tag on SDS-PAGE under reducing (R) condition.


The gel was stained with Coomassie Blue. The purity of the protein is greater than 92% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

Discounts, Gifts,
and more!



» www.acrobiosystems.com



Immobilized Human CEACAM-1, His Tag (Cat. No. CE1-H5220) at 10 μ g/mL (100 μ L/well) can bind Biotinylated Human CEACAM-8, His,Avitag (Cat. No. CE8-H82E9) with a linear range of 0.039-1.25 μ g/mL (QC tested).

Background

Carcinoembryonic antigen-related cell adhesion molecule 1 (CEACAM1) is also known as Biliary glycoprotein 1 (BGP1), CD66a, which belongs to the immunoglobulin superfamily or CEA family. CEACAM1 /CD66a contains three Ig-like C2-type (immunoglobulin-like) domains and one Ig-like V-type (immunoglobulin-like) domain. CEACAM1 /CD66a was described as an adhesion molecule mediating cell adhesion via both homophilic and heterophilic manners, and was detected on leukocytes, epithelia, and endothelia. Studies have revealed that CEACAM1 / BGP-1 performs actions in multiple cellular processes including tissue differentiation, angiogenesis, apoptosis, metastasis, as well as the modulation of innate and adaptive immune responses.

Discounts, Gifts,
and more!



» www.acrobiosystems.com