

Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

➤ *Limited Use & License Disclosure*

BY USE OF THIS PRODUCT, RESEARCHER AGREES TO BE BOUND BY THE FOLLOWING TERMS OF LIMITED USE OF THIS CELL LINE PRODUCT.

- If the researcher is not willing to accept the terms of limited use of this cell line product, and the product is unused, ACRO will accept return of the unused product.
- Researchers may use this product for research use only, no commercial use is allowed.
"Commercial use" means any and all uses of this product and derivatives by a party for profit or other consideration and may include but is not limited to use in: (1) product manufacture; and (2) to provide a service, information or data; and/or resale of the product or its derivatives, whether or not such product or derivatives are resold for use in research.
- This cell line is neither intended for any animal or human therapeutic purposes nor for any direct human in vivo use. You have no right to share, modify, transfer, distribute, sell, sublicense, or otherwise make the cell line available for use to other researchers, laboratories, research institutions, hospitals, universities, or service organizations.
- ACROBIOSYSTEMS MAKES NO WARRANTIES OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, WITH RESPECT TO THE SUITABILITY OF THE CELL LINE FOR ANY PARTICULAR USE.
- ACROBIOSYSTEMS ACCEPTS NO LIABILITY IN CONNECTION WITH THE HANDLING OR USE OF THE CELL LINE.
- Modifications of the cell line, transfer to a third party, or commercial use of the cell line may require a separate license and additional fees. Please contact order.cn@acrobiosystems.com for further details.

Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

Human IL-11 R alpha (Luc) HEK293 Reporter Cell


Catalog No.	Size
CHEK-ATF052	2 × (1 vial contains ~5×10 ⁶ cells)

• Description

The Human IL-11 R alpha (Luc) HEK293 Reporter Cell was engineered to express STAT3 signaling response element. When stimulated with human IL-11 protein, the IL-11/ IL-11 R alpha interaction drives STAT3-mediated luminescence. Inhibition of IL-11 binding to IL-11 R alpha by either anti-IL-11 or anti-IL-11 R alpha antibodies results in a decrease in luminescence.

• Application

- Screen for anti-human IL-11 or anti-human IL-11 R alpha neutralizing antibody.



Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

• Cell Line Profile

Cell line	Human IL-11 R alpha (Luc) HEK293 Reporter Cell
Host Cell	HEK293
Property	Adherent
Complete Growth Medium	DMEM + 10% FBS
Selection Marker	Hygromycin B (40 µg/mL)
Incubation	37°C with 5% CO ₂
Doubling Time	22-24 hours
Transduction Technique	Lentivirus

• Materials Required for Cell Culture

- DMEM Medium (BasalMedia, Cat. No. L120KJ)

Note: If you are unable to obtain the specified DMEM medium (BasalMedia, Cat. No. L120KJ) in China, you may use an alternative DMEM medium (Gibco, Cat. No. 11965-092) or another suitable medium for culturing.

- Fetal bovine serum (CellMax, Cat.No.SA211.02)
- Hygromycin B (Invitrogen, Cat. No. 10687010)

Note: For selection antibiotics, we highly recommend using the specified brand. The activity of antibiotics may vary between manufacturers, so if you choose to use a different brand, it is essential to validate whether the concentration recommended in the culture medium is suitable. Regardless of the brand used, we recommend maintaining a backup culture without selection antibiotics to avoid potential cell loss due to inappropriate antibiotic concentration.

- 0.25% Trypsin-EDTA (1X), Phenol Red (Gibco, Cat. No. 25200-056)
- Penicillin-Streptomycin (Gibco, Cat. No. 15140-122)
- Phosphate Buffered Saline (1X) (HyClone, Cat. No. SH30256.01)
- Complete Growth Medium: DMEM + 10% FBS, 1%P/S
- Culture Medium: DMEM + 10% FBS, Hygromycin B (40 µg/mL), 1%P/S
- Freeze Medium: 90% FBS, 10% (V/V) DMSO
- T-75 Culture flask (Corning, Cat. No. 430641)
- Cryogenic storage vials (SARSTEDT, Cat. No. 72.379.007)
- Thermostat water bath
- Centrifuge (Cence, Model: L550)
- Cell counter (MONWEI, Model: SmartCell200A Plus)
- CO₂ Incubator (Thermo, Model: 3111)

Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

• *Recovery*

1. Thaw the vial by gently agitating it in a 37°C water bath. To minimize the risk of contamination, ensure the cap remains out of the water. Thawing should be completed quickly, typically within 3-5 minutes.
2. After thawing, promptly remove the vial from the water bath and decontaminate it by spraying with 70% ethanol. From this point onward, all operations must be performed under strict aseptic conditions.
3. Transfer the contents of the vial to a centrifuge tube containing 4.0 mL of complete growth medium. Centrifuge at approximately 1000 rpm for 5 minutes.
4. Resuspend the cell pellet with 5 mL **complete growth medium** and transfer the cell suspension into a T-75 flask containing 10-15 mL of pre-warmed **complete growth medium**.
5. Incubate at 37°C with 5% CO₂ incubator until the cells are ready to be split.

• *Subculture*

1. Cell viability may be low after thawing, and full recovery may take up to a week. Monitor the cells daily until the culture reaches 80-90% confluency. At this point, remove and discard the spent medium. Avoid allowing the cells to become over-confluent to ensure optimal cell health.
2. Wash the cells once with sterile PBS. Avoid adding PBS directly onto the cell surface.
3. Add 2 mL of 0.25% Trypsin-EDTA to the T-75 flask. Place the flask at 37°C for 2-3 minutes, until 90% of the cells have detached. Monitor under a microscope to avoid over-trypsinization.
4. Add 6.0 to 8.0 mL of **culture medium** using a pipette and gently rinse the cells from the surface of the T-75 flask. Gently pipette up and down several times to achieve a single cell suspension without cell clumps.
5. Transfer appropriate aliquots of the cell suspension to a new T-75 flask. A subcultivation ratio of 1:4 to 1:8 is recommended. Adjust the ratio based on your specific culture system.
6. Incubate at 37°C with 5% CO₂ incubator.
7. When the cell culture reaches 80-90% confluency, proceed to the next subculture. Avoid over-confluency, as this may negatively impact cell performance in subsequent passages.

Note: After recovery, maintain the cells for 1-2 passages in the complete growth medium not containing the selection marker, if the cells are in good condition, transition to the culture medium containing the selection marker during subculturing.

Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

• *Cryopreservation*

1. When the cell culture reaches 80-90% confluency, remove and discard the spent medium.
2. Wash the cells once with sterile PBS. Avoid adding PBS directly onto the cell surface.
3. Add 2 mL of 0.25% Trypsin-EDTA to the T-75 flask. Place the flask at 37°C for 2-3 minutes, until 90% of the cells have detached. Monitor under a microscope to avoid over-trypsinization.
4. Add 6.0 to 8.0 mL of complete growth medium using a pipette and gently rinse the cells from the surface of the T-75 flask. Gently pipette up and down several times to achieve a single cell suspension without cell clumps. Count the viable cells.
5. Transfer the cell suspension to a centrifuge tube. Centrifuge at 1000 rpm for 5 min at room temperature to pellet the cells.
6. After centrifugation, discard the supernatant. Resuspend the cells in ice cold freezing medium to a concentration of 5×10^6 to 1×10^7 cells/mL.
7. Aliquot the cell suspension into cryogenic storage vials. Place the vials in a programmable cooler or an insulated box placed in a -80°C freezer overnight, then transfer to liquid nitrogen storage for long-term storage.

Note: It is recommended to establish a cell bank at the earliest possible passage for long-term use.

• *Storage Condition*

Cells must be received in a frozen state on dry ice and should be transferred to liquid nitrogen or a -80°C freezer immediately upon receipt. If stored in a -80°C freezer, it is recommended to limit the storage period to no more than two weeks. For long-term preservation, transfer the cells to liquid nitrogen is highly recommended.

Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

• Signaling Bioassay




Fig1. Response to human IL-11 protein (RLU). The Human IL-11 R alpha (Luc) HEK293 Reporter Cell was stimulated with serial dilutions of human IL-11 protein (Cat. No. IL1-H5243). The EC₅₀ was approximately 0.12 µg/mL.




Fig2. Response to human IL-11 protein (FOLD). The Human IL-11 R alpha (Luc) HEK293 Reporter Cell was stimulated with serial dilutions of human IL-11 protein (Cat. No. IL1-H5243). The max induction fold was approximately 31.

Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

• Application

Anti-human IL-11 Neutralizing Antibody Screening




Fig3. Inhibition of human IL-11 protein-induced reporter activity by anti-human IL-11 neutralizing antibody. This reporter cell was incubated with serial dilutions of antibodies in the presence of human IL-11 protein (Cat. No. IL1-H5243) with a final concentration of 0.2 µg/mL. The EC50 of anti-human IL-11 neutralizing antibody is approximately 1.047 µg/mL.

Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

• *Related Products*

<u>Products</u>	<u>Cat.No.</u>
Human IL-11 Protein, His Tag	IL1-H5243
Human VEGF R2 (Luc) HEK293 Reporter Cell	CHEK-ATF044
Human TSLPR (Luc) HEK293 Reporter Cell	CHEK-ATF045
NFAT (Luc) Jurkat Reporter Cell	SCJUR-STF046
STAT3 (Luc) HEK293 Reporter Cell	CHEK-ATF047
NF- κ B (Luc) HEK293 Reporter Cell	CHEK-ATF048
Human EGF R (Luc) HEK293 Reporter Cell	CHEK-ATF049
NFAT (Luc) HEK293 Reporter Cell	CHEK-ATF050
Human PD-1/LAG-3 (Luc) Jurkat Reporter	SCJUR-STF063
Human PD-1 (Luc) Jurkat Reporter Cell	SCJUR-STF064
Human LAG-3 (Luc) Jurkat Reporter Cell	SCJUR-STF065
Human CD16a (158V) (Luc) Jurkat Reporter Cell	SCJUR-STF067
Human CD16a (158F) (Luc) Jurkat Reporter Cell	SCJUR-STF068
Human CD32a (131H) (Luc) Jurkat Reporter Cell	SCJUR-STF069
Human CD32a (131R) (Luc) Jurkat Reporter Cell	SCJUR-STF070
Human CD32b (Luc) Jurkat Reporter Cell	SCJUR-STF071
Human CD64 (Luc) Jurkat Reporter Cell	SCJUR-STF072
Human 4-1BB (Luc) HEK293 Reporter Cell	CHEK-ATF073
Human IL-5 R alpha/CD131 (Luc) HEK293 Reporter Cell	CHEK-ATF074
Human IL-21 R (Luc) HEK293 Reporter Cell	CHEK-ATF051
Human IL-11 R alpha (Luc) HEK293 Reporter Cell	CHEK-ATF052
Human IL-4 R alpha/IL-13 R alpha 1 (Luc) HEK293 Reporter Cell	CHEK-ATF075
Human TrkA (Luc) HEK293 Reporter Cell	CHEK-ATF093
Human IL-31 RA/OSMR (Luc) HEK293 Reporter Cell	CHEK-ATF094
Human IL-10 R alpha/IL-10 R beta (Luc) HEK293 Reporter Cell	CHEK-ATF095
Human GLP-1R (Luc) HEK293 Reporter Cell	CHEK-ATF096
Human CD40 (Luc) HEK293 Reporter Cell	CHEK-ATF097
Human IL-7 R alpha/CD132 (Luc) HEK293 Reporter Cell	CHEK-ATF099
Human GCGR (Luc) HEK293 Reporter Cell	CHEK-ATF103

Human IL-11 R alpha (Luc) HEK293 Reporter Cell Data Sheet

Products

Cat.No.

Human GIPR (Luc) HEK293 Reporter Cell	CHEK-ATF104
Human HVEM (Luc) HEK293 Reporter Cell	CHEK-ATF105
Human BTLA (Luc) Jurkat Reporter Cell	SCJUR-STF106
Human IGF-1 R (Luc) HEK293 Reporter Cell	CHEK-ATF107
NF-kB (Luc) Jurkat Reporter Cell	SCJUR-STF113
TCF/LEF (Luc) HEK293 Reporter Cell	CHEK-ATF114
Human GLP-2R (Luc) HEK293 Reporter Cell	CHEK-ATF128
Human 5-HT1A (Luc) HEK293 Reporter Cell	CHEK-ATF131
Human RANK (Luc) HEK293 Reporter Cell	CHEK-ATF129
Human NKp46 (Luc) Jurkat Reporter Cell	SCJUR-STF130
Human IL-17 RA/IL-17 RC (Luc) HEK293 Reporter Cell	CHEK-ATF133
ISRE (Luc) HEK293 Reporter Cell	CHEK-ATF134
Human OX40 (Luc) HEK293 Reporter Cell	CHEK-ATF135
Human IL-2 R beta/IL-2 R gamma (Luc) HEK293 Reporter Cell	CHEK-ATF136
Human c-MET (Luc) HEK293 Reporter Cell	CHEK-ATF144
Human TGF-beta R (Luc) HEK293 Reporter Cell	CHEK-ATF145
Human FGF-21 (Luc) HEK293 Reporter Cell	CHEK-ATF163
Human Activin RII (Luc) HEK293 Reporter Cell	CHEK-ATF164
Human IL-23 R/IL-12 R beta 1(Luc) HEK293 Reporter Cell	CHEK-ATF166
Human IL-22 R alpha 1/IL-10 R beta (Luc) HEK293 Reporter Cell	CHEK-ATF167
Human DR3 (TL1A receptor) (Luc) Jurkat Reporter Cell	SCJUR-STF178
Human THRA (Luc) HEK293 Reporter Cell	CHEK-ATF180
Human THRB (Luc) HEK293 Reporter Cell	CHEK-ATF181
Human TSHR (Luc) HEK293 Reporter Cell	CHEK-ATF187
Human BMP (Luc) HEK293 Reporter Cell	CHEK-ATF188
Human PTH1R (Luc) HEK293 Reporter Cell	CHEK-ATF194
Human TACI (Luc) HEK293 Reporter Cell	CHEK-ATF197
Human IL-2 R alpha & IL-2 R beta & IL-2 R gamma (Luc) HEK293 Reporter Cell	CHEK-ATF201
Human IL-1 R1 & IL-1 RAcP (Luc) HEK293 Reporter Cell	CHEK-ATF202