

**Synonym**

COL1A1

Source

Mouse COL1A1 Protein, His Tag (CO1-M5243) is expressed from human 293 cells (HEK293). It contains AA Leu 1218 - Val 1453 (Accession # [P11087](#)).

Predicted N-terminus: His

Molecular Characterization

Poly-his	COL1A1(Leu 1218 - Val 1453) P11087
----------	---------------------------------------

[Other Tags and Version](#) [Biotin & Other Labeled Version](#)

This protein carries a polyhistidine tag at the N-terminus.

The protein has a calculated MW of 28.8 kDa. The protein migrates as 33-37 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

The protein is designed as a trimer.

EndotoxinLess than 0.01 EU per μ g by the LAL method / rFC method.**Purity**

>90% as determined by SDS-PAGE.

FormulationLyophilized from 0.22 μ m filtered solution in 50 mM Tris, 150 mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.**Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.



Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System

- [QMS\(ISO, GMP\)](#)
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE**SEC-MALS**

Mouse COL1A1 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than than 85% and the molecular weight of this protein is around 87-102 kDa verified 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

The purity of Mouse COL1A1 Protein, His Tag (Cat. No. CO1-M5243) is more than 85% and the molecular weight of this protein is around 87-102 kDa verified by SEC-MALS.

Background

Type I collagen is the most abundant structural protein of connective tissues such as skin, bone and tendon. It is synthesized as a procollagen molecule which is characterized by a 300 nm triple helical domain flanked by globular N- and C-terminal propeptides. The triple helical domain contains Gly-Xaa-Yaa triplets where Xaa and Yaa are frequently proline and hydroxyproline, respectively. The non-helical propeptides are removed by procollagen N- and C-proteinase activities so that the mature triple helices can self-assemble into collagen fibrils that provide tensile strength to tissues. Type I collagen is a heterotrimer that consists of two alpha 1(I) chains and one alpha 2(I) chain, although homotrimers consisting of three identical alpha 1(I) chains have also been described .

Discounts, Gifts,
and more!



www.acrobiosystems.com

11/5/2025