Human IgG2 Kappa Isotype Control (mAb, carrier free, MALS verified)

Catalog # DNP-M914



Source

Human IgG2 Kappa Isotype Control (mAb) is a chimeric monoclonal antibody recombinantly expressed from HEK293, which combines the variable region of a mouse monoclonal antibody with Human constant domain.

Isotype

Human IgG2 | Human Kappa

Conjugate

Unconjugated

Antibody Type

Recombinant Monoclonal

Reactivity

Human

Specificity

This product is a specific antibody against DNP.

Application

Application	Recommended Usage
ELISA	0.1-8 ng/mL

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Purification

Protein A purified / Protein G purified

Formulation

Supplied as $0.2\ \mu m$ filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Shipping

This product is supplied and shipped with dry ice, please inquire the shipping


Storage

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- The product MUST be stored at -70°C or lower upon receipt;
- -70°C for 3 months under sterile conditions.


SDS-PAGE



Human IgG2 Kappa Isotype Control (mAb) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS




The purity of Human IgG2 Kappa Isotype Control (mAb) (Cat. No. DNP-M914) is more than 90% and the molecular weight of this protein is around 135-155 kDa verified by SEC-MALS.

Report

Human IgG2 Kappa Isotype Control (mAb, carrier free, MALS verified)

Catalog # DNP-M914





Human IgG2 Kappa Isotype Control (mAb) Conc. (ng/mL)

Immobilized DNP-Albumin Conjugate, Bovine at $10 \,\mu\text{g/mL}$ ($100 \,\mu\text{L/well}$) can bind Human IgG2 Kappa Isotype Control (mAb) (Cat. No. DNP-M914) with a linear range of 0.1-1 ng/mL (QC tested).

Background

A hapten is a small molecule that can elicit an immune response only when conjugated with a large carrier such as a protein. Typical haptens include drugs, urushiol, quinone, steroids, etc. Peptides and non-protein antigens usually need conjugating to a carrier protein (such as BSA (bovine serum albumin) or KLH (keyhole limpet hemocyanin) to become good immunogens). Additionally, haptens should be administered with an adjuvant to ensure a high quality immune response. It is important that the hapten design (preserving greatly the chemical structure and spatial conformation of target compound), selection of the appropriate carrier protein and the conjugation method are key conditions for the desired specificity anti-hapten antibodies. We design anti-hapten antibodies based on the HaptenDB information.

